

# Shifting the Value of Norms: Fast Internet, Premarital Sex, and the Erosion of Female Genital Mutilation\*

Jorge García-Hombrados<sup>1</sup>, Daniel Pérez-Parra<sup>2</sup>, and Riccardo Ciacci<sup>3</sup>

<sup>1</sup>Universidad Autónoma de Madrid; IZA

<sup>2</sup>Érudite, Université Gustave Eiffel

<sup>3</sup>Universidad Pontificia Comillas

February 2, 2026

## Abstract

Harmful norms persist because they fulfill a socially valued function. In many Nigerian communities, female genital mutilation (FGM) is practiced because it is believed to discourage premarital and extra-marital sex. This paper examines the impact of the expansion of fast internet on FGM in Nigeria. Our findings indicate that exposure to fast internet reduces both the prevalence of FGM and support for it. The effect does not appear to be driven by exposure to explicit anti-FGM content online. Instead, we find that the effect of fast internet on FGM is driven by fast internet reducing premarital sex stigma, thereby diminishing the value of the social function FGM serves. These findings provide evidence on how harmful norms evolve as the value of their social function changes, with implications for designing effective interventions.

---

\*We received financial assistance from the Comunidad de Madrid (Grant number SI3/PJI/2021-00414) and from the Spanish Ministry of Science and Technology (Grant number PID2020-117402RJ-I00). The qualitative work conducted for this project was crucial for understanding the context, interpreting the results, and guiding the quantitative analysis. This qualitative work was only possible thanks to the superb efforts of the local team: Hadiza Kande Jidda, Anthony Ekere, Stephen Adibe, and Susan Michael. They opened doors to local communities, facilitated every stage of the fieldwork, and made us fall in love with Nigeria and its peoples. We are grateful to Hanna Bony and Javier Martínez Santos for their superb research assistance. Hanna helped assemble a multi-country dataset and prepare the different datasets used in the main analysis. Javier Martínez built the Twitter database. Lindsey Novak and Tamara McGavock facilitated the processed rainfall shock database. We would also like to thank Marine De Talancé, Catherine Bros, Raúl Sánchez de la Sierra, Dean Yang, Sara Lowes, Rebeca Echavarrí, Eugenia Frezza, Suzanna Khalifa, as well as the participants at the DIAL/Dauphine Université, Universidad de Alcalá de Henares, CSAE Research Workshop, the Leibniz University Hannover, Universidad Autónoma de Madrid seminar series, the III Workshop on Public Policies at the Universitat de Barcelona, the BAEM Conference, the CSAE Conference, II Spanish Workshop in Development Economics, and the International Conference in Development Economics for their useful comments.

# 1 Introduction

Despite widespread awareness campaigns and investments in prevention, harmful practices—such as female genital mutilation (FGM), breast ironing, and other harmful bodily modifications—remain prevalent in many communities, affecting hundreds of millions of women worldwide. But why do these norms persist? As damaging as they are, these norms typically serve an important social function. For example, in many traditional communities in Nigeria, where sex outside marriage is highly stigmatized, FGM is believed to discourage premarital sex.

This paper provides evidence on how harmful norms evolve when the social function they serve loses value within a community. Using difference-in-differences methods and granular data on FGM and 3G availability, we assess how the staggered expansion of 3G in Nigeria—which increases fast internet access and exposes people to other cultures and values through films, music, and social networks—affects the evolution of FGM. We first show that the expansion of 3G substantially increased internet access and significantly reduced the prevalence of FGM. 3G availability in the village at the time of birth decreases the probability of undergoing FGM by 2.3 percentage points—approximately 14% of the mean prevalence in the sample. This effect is driven by rural areas, which have historically been more traditional, less exposed to foreign cultural content before the arrival of 3G, and where FGM is more common. Using a sample of men and women 15 or older, we also find that 3G availability at the time of the survey reduces support for the practice.

We also document that the reduction in the prevalence of FGM is closely linked to the effect of 3G in destigmatizing premarital sex. First, we find strong effects of 3G on sexual behavior, shifting attitudes toward more progressive stances: 3G decreases the reported age at first sex while not affecting the age at marriage, increases the reported number of lifetime sexual partners and extramarital relations, and reduces premarital sex stigma. Second, the effect of 3G on FGM is driven by areas where the practice is justified on the grounds of premarital sex stigma. In areas where FGM is justified as a rite of passage for social acceptance, for hygienic reasons, or on religious grounds, but not for preventing premarital sex, the effect is zero. Likewise, the effect of 3G is null in areas where premarital sex is not stigmatized. Taken together, these results show that by reducing premarital sex stigma—and thus undermining the value of the social function that sustains FGM—3G reduced the practice of this harmful norm.

But fast internet constitutes a multidimensional shock whose effects on FGM could, in principle, operate through various alternative mechanisms beyond the destigmatization of premarital sex. We examine and rule out several potential channels

that could plausibly account for the observed effects. Specifically, we find no evidence that the results are driven by exposure to explicit anti-FGM online content or campaigns, or simply by 3G affecting household economic welfare, reducing religiosity, or improving educational attainment.

We examine the robustness of our results through several empirical exercises. First, we show that the effects are robust to alternative difference-in-differences estimators, specification checks, and estimation samples. Second, we discuss and provide evidence that the main conclusions are not driven by measurement error or by 3G affecting the reporting of FGM rather than the true incidence of the practice. Third, by showing that the effect on FGM is not driven by 2G technology, we show that the effect is linked to fast internet specifically, rather than to other phone functionalities. Fourth, we test and rule out the hypothesis that the effect of 3G is confounded by major policies implemented in Nigeria around the same time, which could have affected FGM or its reporting, or by an increase in the presence of the state resulting from the expansion of 3G. Finally, we show that the results are not simply driven by households more opposed to FGM migrating towards 3G covered areas.

Nigeria provides a highly relevant and ideal setting to examine the effect of fast internet on the evolution of FGM. First, this country is the largest mobile broadband market in Africa, with around 170 million mobile connections in 2019 (Bahia et al., 2024). Second, while the prevalence of FGM in Nigeria is not as high as in other countries such as Somalia, Mali or Guinea, Nigeria has the third largest number of cut women after Egypt and Ethiopia. It is estimated that 10% of all women who have undergone FGM worldwide are Nigerian.<sup>1</sup> Third, unlike many other countries, the vast majority of cuts in Nigeria occur during childhood. This is specially important to define the risk of being cut over the children's first years of life, which is an essential part of the identification strategy used in this paper. Finally, rich granular data from Nigeria enable our proposed analysis. We merge several geocoded individual-level data sources from Nigeria that include information on FGM, support for the practice, and sexual behavior, covering both the period before and after the arrival of 3G in the early 2010s, with yearly-level data on 3G availability at the 1km×1km grid level until 2018. We complement these datasets with focus groups that we implemented in December 2024 to gain better insights about internet use and its perceived effects on social norms in Nigeria.

Our study speaks to different streams of research.

---

<sup>1</sup>See the following link <https://www.fgmcri.org/blog/nigerias-battle-against-female-genital-mutilation-a-hard-look-at-the-numbers-and-the-fight-ahead/>

First, we contribute to the multidisciplinary literature on the evolution of social norms by providing empirical evidence that supports the hypothesis that norms erode when the social functions they fulfill become obsolete. This perspective aligns with theoretical frameworks developed in [Bicchieri \(2005\)](#), [Henrich \(2015\)](#), [Giuliano and Nunn \(2020\)](#), [Henrich \(2020\)](#), and [Henrich and Muthukrishna \(2021\)](#), which highlight how norms are maintained through social learning, coordination dynamics, and their adaptive value to communities. In other words, norms persist and evolve to solve social problems ([Henrich et al., 2001](#); [Henrich and Muthukrishna, 2021](#)). In the same spirit, but using an economic lens, [Fernández \(2025\)](#) reviews seminal economic work and proposes a framework for cultural change based on incentives, showing how cultural practices adapt when the incentives underpinning them shift. Our study shows that FGM tends to be less frequent when premarital sex is normalized, as the value of FGM’s social function—preventing premarital sex—declines and the underlying social incentives for cutting are reduced. More broadly, our results suggest that norm change can be understood as a function of shifts in cultural relevance of the behaviors that norms are designed to regulate, contributing to broader debates about the co-evolution of culture and norms.

Second, our study contributes to the body of evidence on what works to fight FGM. This harmful norm has lasting health consequences ([Jones et al., 1999](#); [Berg et al., 2014](#); [Wagner, 2015](#)). However, despite the substantial resources invested in fighting FGM ([Katz et al., 2021](#)), the practice affects 4 million girls every year ([WHO, 2025](#)). Policies to combat FGM often focus on improving maternal education and women’s bargaining power. While these policies have well-established welfare effects for women, the evidence suggests that they have limited impact on FGM itself ([De Cao and La Mattina, 2019](#)). So, what works to fight this practice? [Novak and Bussberg \(2023\)](#) review the evidence addressing this question and conclude that public policies have rarely been rigorously evaluated, and those that have been show only limited observable impacts.<sup>2</sup> The review emphasizes that fostering changes in FGM is difficult because it requires tackling the cultural and identity roots of the practice, which vary substantially across ethnic groups. Evidence on the effectiveness of such approaches, however, remains scarce ([Corno and La Ferrara, 2019](#); [Congdon Fors et al., 2024](#)). We add to this literature by examining the role of fast internet—arguably one of the most powerful catalysts of cultural change in sub-Saharan Africa in the 21st century. While the expansion of fast internet may have adverse effects on other outcomes, our findings suggest that interventions weakening the value of the social functions that sustain harmful norms can be an

---

<sup>2</sup>See Section 2 for a broader discussion on the effects of anti-FGM policies.

effective strategy to reduce FGM and similar practices.

Third, our study contributes to the economics literature on the determinants and the nature of FGM. [Becker \(2024\)](#) argues that infibulation—the most severe form of FGM—emerged in pastoralist societies as a mechanism to control female sexuality and limit extramarital sexual relations during the prolonged absences of husbands and [La Ferrara et al. \(2020\)](#) show that the Red Sea slave trade increased the prevalence of the practice, which was used to signal sexual chastity among enslaved women. [Poyker \(2023\)](#) shows how regime stability can facilitate the abandonment of FGM, and [Gulesci et al. \(2023\)](#) use data from Somalia to document how FGM practices can shift rapidly from more severe to less severe forms, consistent with the idea of incremental norm change through “stepping stones”. [Novak \(2020\)](#), [Bellemare et al. \(2015\)](#), and [Efferson et al. \(2015\)](#) examine whether FGM operates as a social coordination norm in different sub-Saharan African countries. They find no evidence of a single tipping point and show that much of the variation in prevalence reflects household-level characteristics. Our study contributes to this body of literature by providing empirical evidence supporting the anthropological hypothesis that links FGM to cultural beliefs and stigmas around premarital sex ([Koso-Thomas, 1987](#); [Shell-Duncan and Hernlund, 2001](#); [Ahmadu, 2001](#)).

Finally, our study contributes to the growing literature on the transmission of norms and culture. Previous literature has highlighted the role of TV shows, social media or migrants in the spread of norms and the formation of values in developing countries ([Jensen and Oster, 2009](#); [La Ferrara et al., 2012](#); [Diabate and Mesplé-Somps, 2019](#); [Ahmed et al., 2025](#)). More specifically, a handful of studies have explored the role that new communication technologies and internet have on social norms and behaviors. [Zhuravskaya et al. \(2020\)](#) and [Guriev et al. \(2020\)](#) delve into the influence of 3G networks on political outcomes, [Manacorda and Tesei \(2020\)](#) reveals the mobilization and protest-facilitating role of mobile phone coverage (2G) in Africa, and [Frezza \(2023\)](#) uncovers adverse effects on women’s welfare and justification of intimate partner violence in West Africa stemming from the arrival of both terrestrial and submarine cables.<sup>3</sup> We show that fast internet acts as a catalyst for broader cultural change, with effects on harmful norms such as FGM and on sexual behaviors.

The rest of this paper is structured as follows. Section 2 characterizes the practice of FGM, presenting stylized facts, and summarizing the evidence on the effectiveness of anti-FGM policies. Section 3 describes the expansion of fast internet in the anal-

---

<sup>3</sup>The effects of new communication technologies on economic outcomes are discussed in [Hjort and Poulsen \(2019\)](#) and [Bahia et al. \(2024\)](#).

ysis and why it might have changed FGM. Sections 4 and 5 introduce the empirical strategy and the data used in the main analyses. Section 6 presents the results of the expansion of 3G on the prevalence of FGM. Section 7 shows that the main effects on FGM are driven by 3G-induced changes in sexual behavior. Section 8 examines and rules out alternative hypotheses that could explain the effect of 3G on FGM. Section 9 discusses the results of the qualitative analysis on which types of internet content may be driving the effect of 3G on sexual behaviors. Section 10 concludes.

## 2 Female genital mutilation: How widespread is it, why is it practiced, and what works against it?

Defined as the total or partial removal of female genitalia for reasons unrelated to health, the practice of FGM affects around 230 million women worldwide (UNICEF, 2024). Figure A1 in Appendix A shows the prevalence across various countries where this practice is existent among women aged 15–49. The figure also displays how prevalence varies across generations, suggesting that while the practice is decreasing in some countries such as Burkina Faso or Sierra Leone, it remains largely unchanged in others such as Somalia or Guinea. The WHO estimates that 4 million girls are still at risk of FGM every year (WHO, 2025).

But why is FGM practiced? A multidisciplinary body of literature has examined its multifaceted meanings and causes, which vary across different societies. Gruenbaum (2001) and Koso-Thomas (1987) highlight that in Sudan and Sierra Leone, FGM signals adherence to societal norms, rites of passage, and the preservation of tradition. Within the Kono community in Sierra Leone, Ahmadu (2001) documents that FGM can be also seen as a marker of female empowerment and identity. Mandara (2001) argues that in Nigeria, FGM is believed to reduce women’s sexual desire, thereby encouraging loyalty and discouraging premarital sex. Khalifa (2022) and García-Hombrados and Salgado (2022) show that in Egypt and Senegal, FGM facilitates marriage and may lead to a higher bride price. Overall, the literature documents that within contexts where FGM is practiced, it is often deeply linked to women’s identity (Shell-Duncan and Hernlund, 2001), reflecting characteristics that are socially valued and esteemed, including femininity, purity, sexual loyalty to the husband, maturity, and stability.

A common belief among policymakers and the general public associates FGM with a lack of education and women’s bargaining power. While there is a well-

established negative correlation between maternal education and the likelihood of FGM, [De Cao and La Mattina \(2019\)](#) show that maternal education has no causal effect on either the prevalence of FGM or its support. Moreover, DHS data reveal that in most countries, support for FGM is similar among women and men.<sup>4</sup> Considering that women are already the primary decision-makers in most cutting decisions ([Novak and Bussberg, 2023](#)), it is unclear whether improving mothers' bargaining power would lead to a significant reduction in FGM.<sup>5</sup>

So, what works to reduce FGM? [Novak and Bussberg \(2023\)](#) review the evidence on which policies are effective in combating FGM, concluding that there is very little rigorous evidence on what works. The largest effects are found for interventions and shocks that target the cultural roots of FGM, such as the adoption of alternative rites of passage ([Corno and La Ferrara, 2019](#)) or spillover effects from returned migrants ([Diabate and Mesplé-Somps, 2019](#)). In line with the relevance of targeting the cultural roots of the practice, [Congdon Fors et al. \(2024\)](#) shows that Christian missions in sub-Saharan Africa decreased the current prevalence of FGM, and that this effect was not driven by better education or economic development. The evidence on the effects of legal bans is more mixed: some studies suggest positive impacts in Burkina Faso and Senegal ([Crisman et al., 2016](#); [García-Hombrados and Salgado, 2022](#)), while others find no effects or even unintended negative consequences in terms of the age at cutting ([Camilotti, 2015](#); [Cetorelli et al., 2020](#)). Finally, interventions focused on information provision, campaigns, or advocacy can reduce the prevalence ([Vogt et al., 2016](#); [Corno and La Ferrara, 2019](#); [Khalifa, 2022](#)) or the severity of the practice ([Gulesci et al., 2020](#)), though they can also provoke backlash ([Gruenbaum, 2001](#)).

## 2.1 FGM in Nigeria

The most recent round of the Nigerian DHS, conducted in 2018, indicates that approximately 20% of women aged 15–49 have undergone FGM.<sup>6</sup> [Figure 1](#) shows the prevalence of FGM by year of birth in Nigeria. Despite a significant decline among women born in the last decades of the 20th century, prevalence stabilized in the late 1990s and experienced a slight increase in the following years, before decreasing again for cohorts born in the 2010s, coinciding with the expansion of 3G.

---

<sup>4</sup>See [Figure A2](#) in [Appendix A](#).

<sup>5</sup>An exception is provided by [Harari \(2019\)](#), who shows that in Kenya, the expansion of women's inheritance rights reduced the prevalence of FGM.

<sup>6</sup>While a MICS survey was conducted in Nigeria in 2021, we use the 2018 Nigerian DHS to present descriptive evidence on FGM, as this is the last year for which information on 3G is available and therefore the last year included in our study period. The prevalence of FGM in the 2021 MICS among women aged 15–49 is 16%.

The survey also shows that most procedures involve partial or total removal of the clitoral glans, but not infibulation, and are carried out by traditional circumcisers. Unlike in other settings, our DHS data for Nigeria shows that more than 80% of cuts occur within the first year of life, and more than 95% before age 3.<sup>7</sup> FGM is deeply rooted in the traditions of Nigeria’s most populous ethnic groups, including the Yoruba, Igbo, and Hausa. Figure 2 shows that while FGM is present throughout the country, prevalence is higher in the South and Southeast, with certain focal points in Northern Nigeria.<sup>8</sup> In most communities surveyed in the DHS data, FGM is practiced on less than half of the relevant population, suggesting that in Nigeria it is unlikely to function purely as a social coordination norm and that decisions vary across households within the same communities.

The practice of FGM in Nigeria shares commonalities with other contexts, as it is used to promote behaviors that are socially valued—in the case of Nigeria, virginity and loyalty (Orubuloye et al., 2001). Indeed, across most ethnicities, FGM is widely believed to reduce sexual desire and discourage premarital and extra-marital sex (Mandara, 2001; Orubuloye et al., 2001), behaviors that are strongly stigmatized in many traditional communities. Consistent with this, the latest Nigerian DHS data on perceived benefits of FGM (2008 round) highlight the preservation of virginity as the most commonly cited benefit (see Figure 3). Other non-mutually exclusive benefits of FGM cited by the survey respondents include better marriage prospects, social acceptance, improved hygiene, more sexual pleasure for men, and, less frequently cited, religious approval.<sup>9</sup> While the importance of virginity is sometimes associated with women’s prospects in the marriage market, our qualitative analysis suggests that, in Nigeria, virginity carries a broader social meaning. Beyond its potential implications for marriage, remaining a virgin until marriage is often seen as a marker of moral integrity and family honor, conferring higher social status and greater respect from peers, relatives, and even ancestors (Ademiluka, 2023; Taiwo et al., 2023).

---

<sup>7</sup>For example, in some traditional Igbo communities, FGM was sometimes performed during the naming ceremony, seven days after birth.

<sup>8</sup>The prevalence of FGM at the county level in 2008, before the deployment of 3G, is shown in Figure A3 in Appendix A. DHS surveys are not representative at the county level, but rather at the regional level for women aged 15–49. The prevalence for this age group at the regional level in 2008 and 2018 is presented in Figure A4 in Appendix A.

<sup>9</sup>The link between religion and FGM is discussed in detail in Section 6.3.

### 3 The spread of 3G in Nigeria and how it might affect FGM

To explore the effect of fast internet on FGM in Nigeria, we focus on the expansion of the third and fourth generation of mobile telecommunications, known as 3G and 4G networks. 3G and 4G technology provides enhanced mobile communication capabilities, including fast data transmission and expanded multimedia capabilities, allowing for fast internet access from mobile phones. While we consider both 3G and 4G for our analysis, 4G was first introduced in Nigeria at the very end of our study period and typically in areas that previously already had 3G. For simplicity, we use the term 3G throughout the paper.

This technology operates through cell towers or base stations that communicate with user terminals in their vicinity using electromagnetic signals to connect these terminals to the core network and provide fast internet. The reach of cell towers can extend up to 15 km, depending on topography, buildings, transmission power, and weather conditions. Introduced first in the main Nigerian cities, 3G networks gradually extended nationwide during the 2010's. By the end of 2019, Nigeria had 170 million mobile connections (60% of which used 3G or 4G mobile broadband technologies), which makes Nigeria the largest mobile broadband in Africa ([Bahia et al., 2024](#)). Figure 4 shows the expansion of 3G availability throughout the country.

In Sub-Saharan Africa, where cable connections, DSL and fiber optic infrastructures are limited and costly (less than 0.5 subscriptions per 100 inhabitants ([World Bank, 2024](#))), 3G mobile networks, accessed via smartphones, serve as the predominant means of internet access. Using data from the Nigerian General Household Survey (NGHS), Figure A5 in Appendix A shows how internet access started increasing from 2012, when the rollout of 3G networks began, with mobile phones becoming the main means of access (see Figure A6 in Appendix A). In Section 6, we use the NGHS to document large causal effects of 3G availability on internet access, particularly among young women.

We hypothesize that fast internet can erode the practice of FGM through several mechanisms. We provide a brief overview of the mechanisms here, with a fuller discussion in Sections 7 and 8. First, the internet facilitates the dissemination of explicit anti-FGM content and campaigns, granting individuals the opportunity to access such materials. While pro-FGM content is also available online, it is far less common ([Babbs et al., 2023](#)). A detailed discussion of these online campaigns and efforts is provided in Section 8.1. But most of the internet content accessed by individuals, however, is unlikely to be about FGM. Could non-explicitly FGM

content decrease FGM? The internet may also expose individuals in traditional communities to cultural content and role models that portray female identities in which premarital sex is more common than what is typically seen in traditional communities in Nigeria. Considering that FGM is mainly practiced in traditional communities because it is believed to discourage women from engaging in sex outside marriage, the need for such a practice may decrease if prevailing ideas about women’s sexual behaviors change due to exposure to more progressive ideals and role models on the internet (e.g. music, YouTube videos, movies, social media, etc.). On the other hand, exposure to different cultures, behaviors, or simply anti-FGM content could also trigger a cultural backlash (Blumenstock et al., 2022; Gruenbaum, 2001), raising questions about the direction of the effect—an issue that remains an open empirical question. Finally, fast internet is also a broad socioeconomic shock that may have multidimensional consequences, some of which may ultimately affect the prevalence of FGM. In Section 8, we discuss and examine the extent to which the effect of 3G on FGM could be driven by improvements in economic conditions, a decrease in religiosity, or by an increase in education.

## 4 Empirical strategy

To assess the impact of fast internet on FGM, we leverage information obtained from the daughters of women participating in the 2018 and 2013 rounds of the Nigerian Demographic Health Survey (DHS) and in the 2021 and 2016 rounds of the Nigerian Multiple Indicator Cluster Survey (MICS) that were born between 2007 and 2018, the years for which we have granular information about 3G availability. The information is collected from the birth registry module which includes information on FGM status for every daughter of women aged 15-49.<sup>10</sup>

Our empirical approach relies on variation in the availability of 3G network at the time of birth in the place of residence.<sup>11</sup> Because FGM in Nigeria occurs mainly within the first year of life, those girls born after the introduction of this technology in the village/ward of residence are considered exposed to 3G while, those born before 3G network is available in the location or in locations with no 3G network available during the study period, are considered unexposed. Formally, we estimate the following difference-in-differences model:

---

<sup>10</sup>While the female module collected self-reported information on FGM for women aged 15 and above, all girls born between 2007 and 2018 were 14 or younger at the time of the survey. Consequently, the FGM status for all girls in the analytical sample is reported by their mothers.

<sup>11</sup>The DHS and MICS datasets do not include the coordinates for the place of birth. We show in Section 6.3 that the results are robust to the exclusion of migrant households.

$$FGM_{ibv} = \delta_0 + \delta_1 3G\ available_{bv} + \theta_b + \gamma_v + u_{ibv} \quad (1)$$

where  $FGM_{ibv}$  is equal to 1 if girl  $i$ , born in year  $b$  and living in village  $v$  is cut and 0 otherwise.  $3G\ available_{bv}$  indicates whether village/ward  $v$  in year  $b$  is covered by 3G network or higher generation.  $\delta_1$  is the parameter of main interest and it measures the effect of 3G network availability at the time of birth on the probability of FGM.  $\theta_b$  and  $\gamma_v$  are year-of-birth and village fixed effects to control for time-invariant village-level characteristics and birth-year specific events. Standard errors are clustered at the village/ward level (the survey cluster).

Because the introduction of 3G throughout Nigeria is staggered, the standard OLS model may lead to the negative weights problem identified by [Goodman-Bacon \(2021\)](#). We decompose the weights and find that less than 1% of the weights used in the OLS estimation are negative, validating the use of the canonical TWFE model. As a robustness check, we also estimate the average treatment effects using the difference-in-differences for staggered treatment adoption methods developed in [Sun and Abraham \(2021\)](#), [Callaway and Sant’Anna \(2021\)](#), and [Chaisemartin and D’Haultfœuille \(2020\)](#) finding nearly identical estimates. The deployment of 3G networks over time was unlikely to be random and was more likely driven by the costs and potential returns of expansion.<sup>12</sup> However, the validity of our difference-in-differences approach does not require the expansion to be random. Instead, the key identifying assumption in our strategy is the parallel trends condition: In absence of 3G network, the evolution of the prevalence of FGM would be similar across cohorts exposed and unexposed to 3G. We examine the plausibility of this assumption by estimating event-study specifications.

We also assess the effect of 3G network availability on other outcome variables, mainly, the support for FGM and sexual behavioral outcomes. Information on these variables at the time of the survey is included for women aged 15-49, and for men aged 15-59 in DHS rounds 2003, 2008, 2013 and 2018.<sup>13</sup> Unlike in the analysis of the FGM prevalence, exposure to 3G is not defined based on the availability of 3G network at the time of birth, but rather at the time of the survey, and the population of interest are not the young daughters but rather men and women who are at least 15 years at the time of the survey. As the DHS data are repeated cross-sections and the survey clusters are different in every round, village fixed effects cannot be estimated as this would remove any time variation. We circumvent this limitation

---

<sup>12</sup>These, in turn, may depend on factors such as topography, population density, and grid availability, among others.

<sup>13</sup>Not all the variables are available in every survey round for both men and women.

by converting our treatment variable of interest, 3G exposure, into a variable representing the proportion of the population covered by 3G networks in each county (Local Government Area, LGA), the vast majority of which are surveyed in every DHS round.<sup>14</sup> Defining the treatment at the county level enables us to incorporate county fixed effects and cluster observations by county, allowing *de facto* the estimation of difference-in-differences models. We estimate the following specification:

$$\text{SupportFGM}_{itl} = \delta_0 + \delta_1 \text{Sh population with 3G available}_{itl} + \mu_t + \nu_l + X_{itl} + u_{itl} \quad (2)$$

where  $\text{SupportFGM}_{itl}$  indicates the outcome of interest for individual  $i$ , living in county  $l$  interviewed at time  $t$ ,  $\text{Sh population with 3G available}_{itl}$  is the share of the population in county  $l$  at time  $t$  that has 3G network available.  $\mu_t$  and  $\nu_l$  are year-of-interview (survey) and county (LGA) fixed effects. Finally,  $u$  is the error term. Standard errors are clustered at county level. Unlike in the analysis of FGM, the results of the Goodman-Bacon decomposition calls for caution about the potential problems of negative weights for most outcomes. Thus, we also use the staggered difference-in-differences estimator presented in [Chaisemartin and D’Haultfoeuille \(2020\)](#). The latter approach is the only difference-in-differences estimator for staggered adoption that allows for the use of continuous treatment variables.

To test the feasibility of the parallel trends condition in this specification with a continuous treatment indicator and staggered expansion, we estimate the placebo analysis proposed in [Chaisemartin and D’Haultfoeuille \(2020\)](#). Additionally, we test whether changes in the outcome of interest between the 2003 and 2008 surveys—prior to the start of 3G deployment in Nigeria—are correlated with the degree of 3G availability in the county in 2018.

As a robustness check, we define exposure to 3G at the time of the survey at the individual level to estimate the association between 3G network availability at the time of the survey and these outcomes in [Appendix B](#), showing consistent estimates.

## 5 Data

This research combines quantitative data from multiple sources and qualitative data collected by the authors in the field. Below, we provide information on each of the databases used.

---

<sup>14</sup>There are 774 Local Government Areas (LGAs) in Nigeria. These are equivalent to counties in many other countries and represent the administrative level above wards. LGAs are responsible for local governance and the provision of basic public services.

**Nigerian Demographic and Health Surveys (DHS):** We use information from four rounds of the Nigerian DHS conducted in 2003, 2008, 2013, and 2018. DHS data include repeated cross-sections of individual-level information from a nationally representative sample of women aged 15-49 in Nigeria. The questionnaire provides information on demographic, socioeconomic, and health characteristics, including self-reported information on their FGM status, support for FGM, beliefs about FGM, and sexual behavioral outcomes. Moreover, in the 2018 and 2013 rounds, the survey asked these women about the FGM status for each living daughter of the interviewed women, which is used in the main analysis. Information on the perceived benefits of FGM is only included in the 2003 and 2008 surveys. The survey also includes a male module that collect information on a nationally representative sample of men aged 15-59. The men sample is smaller since only one third of the interviewed households were selected for the male module. Moreover, the male module only includes information on support for FGM in the 2003, 2008, and 2013 rounds. All the DHS rounds considered include information on the latitude and longitude of the survey cluster (village for rural areas and wards for urban areas).<sup>15</sup>

During the survey implementation and in particular the FGM module, enumerators are instructed to ensure privacy, though achieving complete seclusion from other family members may not always be feasible. The words and methods employed by enumerators are specifically designed and trained to avoid underreporting due to social desirability bias and other causes (USAID, 2011). Nonetheless, some degree of underreporting of the FGM status of daughters is expected, and in Section 6.3 we explore the different forms in which underreporting might affect the main estimates and show that our conclusions are not driven by it.

**Nigerian Multiple Indicator Cluster Surveys (MICS)** We use geolocated information from the 2016 and 2021 rounds of the Nigerian MICS on the FGM status of each living daughter of a nationally representative sample of women aged 15–49. This complements the DHS data from the 2013 and 2018 rounds, which are used to analyze the effect of 3G availability at the time of birth on FGM. The MICS data collection process follows procedures and protocols similar to those of the DHS.

**3G data:** We leverage yearly data on 3G and 4G mobile networks coverage in Nigeria available for the period 2007 and 2018.<sup>16</sup> This dataset includes coverage

---

<sup>15</sup>To avoid personal identification, survey clusters are randomly displaced by up to 2 km in urban areas and up to 5 km in rural areas, which may introduce measurement error in the variable on 3G availability. Since this measurement error in the treatment variable is random, our estimates might be thought to be a lower bound for the effect of 3G networks on the outcomes of interest.

<sup>16</sup>The version of the data to which the authors had access does not include information on signal quality. Existing information from OpenSignal indicates that, in 2019, the average mobile internet speed in Nigeria was 9.87 Mbps, one of the highest in sub-Saharan Africa and slightly above other

data at the 1x1-kilometer grid level provided by the Global System for Mobile Communications Association (GSMA).<sup>17</sup> The available information allows therefore to determine whether the individuals interviewed in the different geolocated surveys have 3G or 4G network available at a certain point in time. 4G was first introduced at the very end of our study period and typically in areas that previously already had 3G. Therefore, most of the variation used in the study in exposure to fast internet arises therefore with the deployment of 3G. Thus, while we consider both 3G and 4G to define the treatment variable, we use the term 3G throughout the paper for simplicity.

While we provide evidence that 3G increases internet usage using the Nigerian General Household Survey, our main analysis focuses on 3G (fast internet) availability rather than internet usage for two reasons. First, the DHS dataset, which is the primary dataset including information on FGM outcomes, does not include information on internet usage.<sup>18</sup> Second, 3G availability in the community may also affect households in the community who do not have direct access to the internet. This can occur if, for example, internet users share information and content with non-users in their community, or through the assimilation of behaviors exhibited by internet users in their community.

**Other quantitative datasets:** Following [Gurieva et al. \(2020\)](#), we merge the NASA map of population density at the 1x1 km level<sup>19</sup> with the GSMA dataset on 3G availability to calculate the proportion of population in each Nigerian county every year with 3G network available. This variable serves as the main indicator of treatment exposure in the analysis of the effect of 3G on support for FGM and sexual behavior outcomes. To validate the implicit assumption that 3G network availability increases internet use relevantly, we use information from rounds 2010, 2012, 2015, and 2018 of the Nigerian General Household Survey (NGHS), which collects information on internet use for a nationally representative sample of households. Finally, we use yearly rainfall data collected by the University of Delaware at the grid level for the period 1900-2018. Following [McGavock and Novak \(2023\)](#), we use this dataset to assess the effect of rainfall shocks on FGM in Nigeria, testing whether the estimated impacts of 3G are simply driven by an improvement in economic welfare.

Table 1 shows the descriptive statistics for the data used in the two main analyses.

---

major countries in the Global South, such as India.

<sup>17</sup>The GSM Association serves as the representative body for the collective interests of mobile network operators worldwide.

<sup>18</sup>Only the 2018 round includes this information, which is provided only at the time of the survey and does not allow for use in the retrospective analysis conducted in the main analysis.

<sup>19</sup>The data can be downloaded using this link: <https://neo.gsfc.nasa.gov>

In Panel A, we present descriptive statistics for the sample of females included in the main analysis of the effect of 3G at the time of birth on FGM. The sample consists of daughters of women surveyed in DHS 2013 and 2018, and MICS 2016 and 2021, restricted to those daughters born between 2007 and 2018, the years for which granular data on 3G availability are available. This sample includes cohorts of females born up to five years before the deployment of 3G networks in Nigeria began. The prevalence of FGM in the analytical sample is 16.9%, with 10% of them born in an area covered by 3G at the time of birth. Their ages at the time of the survey range from 0 to 14, with an average age of 4.9 years.

In Panel B, we present descriptive statistics for the sample of males and females included in the analysis of the effect of 3G at the time of the survey on support for FGM, sexual behavior, and other outcomes. The sample consists of men aged 15–59 and women aged 15–49 surveyed in the DHS rounds of 2003, 2008, 2013, and 2018. The number of observations varies substantially across variables and gender because the FGM module was applied only to a subsample of surveyed households, because some questions only targeted ever married women, and because while all women in surveyed households were interviewed, only one-third of the eligible men in these household were selected for the survey. Indeed, more than 70% of the analytical sample is female. The data also show that surveyed individuals are on average around 30 years old, and approximately 30% of respondents believe that the practice of FGM should continue, with the same share for men and women. Age at marriage is low, particularly among women, who have a mean age at first marriage below 18. Extra-marital sex is also uncommon among women, with the average number of lifetime sexual partners excluding husbands being 0.6, while this number is substantially higher for men, at 2.4. Further details about the data sources for the outcome variables used in the paper are provided in Table A1 in Appendix A.

**Qualitative data:** We complement the quantitative analysis with focus groups conducted in the field. In December 2024, two of the authors, in collaboration with local translators, conducted 10 focus groups in Igbo, Yoruba, Hausa, Efok, Ekoi and Ibibio rural communities throughout Nigeria. In them, we collect information on internet use and its perceived effects on social norms, particularly sexual behavior and premarital sex stigma. Details on data collection and analysis are provided in Appendix C.

## 6 Results

### 6.1 The effect of 3G network on internet use

Before presenting the main estimates of the effect of 3G on FGM, we start by examining the effect of 3G availability on internet access. Because information on internet use is not included in Nigerian DHS,<sup>20</sup> we use data from rounds 2010, 2012, 2015, and 2018 of the NGHS. Because the mentioned database surveys the same villages over time, we can estimate a slight variation of equation 2 using individual level internet access as the dependent variable, village-level fixed effects, and defining treatment exposure as a dichotomous variable indicating whether the village had 3G network coverage in the survey year.<sup>21</sup>

The results of this analysis are depicted in Figure 5 using the Sun and Abraham estimation procedure and in Figure A7 in Appendix A using the TWFE.<sup>22</sup> The results of our preferred specification show that the availability of 3G network increases the probability of internet use by approximately 20 percentage points for both men (an increase of 250% relative to pre-3G levels) and women (an increase of 405% relative to pre-3G levels). The effect on internet access is immediate and particularly pronounced among women under 30, increasing by 31 percentage points. These results are not surprising, since 3G is the primary technology for internet access in Nigeria (see Figure A6 in Appendix A), and use of internet in 3G covered areas in Nigeria is large, particularly among young people.<sup>23</sup> On the other hand, infrastructure for cable connections, DSL, and optic fiber is expensive and deficient outside major cities in sub-Saharan Africa (World Bank, 2024).

### 6.2 The effect of 3G network on FGM

The primary focus of the study centers on examining the impact of 3G network availability on the prevalence of FGM. The results of this analysis are reported in Table 2. The effect of 3G availability on the prevalence of FGM ranged from 2.1 to 2.3 percentage points, depending on the difference-in-differences estimator used.

---

<sup>20</sup>Only 2018 DHS round includes information on the use of internet, and this is not asked retrospectively.

<sup>21</sup>While the survey was initially designed as a household panel, the attrition rate was high and many households were replaced. Thus, we set the fixed effect level at the village level. The results were however nearly identical with the fixed effects set at the household level.

<sup>22</sup>Negative weights account for 16–17% of the weights used in the TWFE specification, suggesting that staggered difference-in-differences estimators, such as the one developed by Sun and Abraham (2021), may provide more reliable estimates in this analysis than TWFE estimates.

<sup>23</sup>Round 2018 of the NGHS shows that nearly 55% (66%) of women (men) below 30 living in 3G covered areas declared having access to internet. The % of individuals among the full population in 3G covered areas that access internet is smaller: 39% for women, and 54% for men.

While small in percentage points, these magnitudes correspond to nearly 12% and 14% of the sample mean. The results of the event study reported in Figure 6 reveal that the effects are not driven by pre-existing differential trends.<sup>24</sup> Moreover, the estimates show that the effect already seems to operate for women born in the year when 3G was deployed ( $t_0$  in Figure 6). Although social norms are often thought to be slow-moving, cultural change can sometimes occur quite rapidly. A growing body of research shows that exposure to diverse cultural content or new forms of social interaction has, in some contexts, triggered abrupt shifts in prevailing norms and behaviors, rather than the gradual transitions typically expected (Jensen and Oster, 2009; Chong and Ferrara, 2009; La Ferrara et al., 2012; Fernández, 2025; Ahmed et al., 2025).

We then explore the heterogeneous effects of 3G network availability along three dimensions: areas where FGM is the dominant norm versus those where it is not, rural versus urban areas, and Muslim versus non-Muslim populations. To do so, we re-estimate equation 1 including the relevant characteristic that defines each heterogeneity dimension, as well as its interaction with 3G network availability. The coefficient on this interaction captures the differential effect of 3G availability on FGM across that particular dimension of heterogeneity. The results reported in column 1 of Table 3 show nuanced evidence of the heterogeneous effects of 3G availability by whether FGM is the dominant rule. While the magnitude of the effect is approximately 68% larger in survey clusters where the prevalence of FGM is more than 50%,<sup>25</sup> the effects in survey clusters where FGM is the prevalent norm are not statistically different from those in areas where it is not. This result should be interpreted with caution because, in only about 20% of these clusters, women subjected to FGM are in the majority. The results for the comparison between urban and rural areas are reported in column 2. The estimates show that the effect of 3G availability on FGM is fully driven by rural areas. The effect on urban areas is small and largely insignificant. These results are consistent with the hypothesis that the effect of exposure to fast internet is stronger in communities with lower levels of exposure to other cultures before the deployment of 3G network. Finally, the results reported in column 3 show that the coefficient on the interaction between 3G availability and being Muslim is negative, consistent with a larger effect for Muslims, who are the majority in the north of the country. However, once again,

---

<sup>24</sup>Figure A8 in Appendix A show that the effects are similar when estimated using the event study method developed in Sun and Abraham (2021).

<sup>25</sup>The cluster prevalence in this analysis is calculated using the sample of women aged 15–49 in the cluster. Women in this age range were at least 9 years old when 3G was first deployed, and their FGM status is therefore unlikely to have been affected by 3G. Moreover, these women are not included in the analytical sample.

this difference is statistically indistinguishable from zero at conventional confidence levels.

### 6.3 Robustness of the results

Using different difference-in-differences estimators, the results reported above show that 3G network coverage at the time of birth reduced the probability of FGM. In this subsection, we explore the robustness of our findings to different empirical exercises and identification threats.

First, we re-estimate specification 1 adding controls for age, access to electricity, TV, and radio. Results are reported in columns 2 and 3 of Table 4, with column 1 replicating the benchmark estimates from Table 2 for ease of comparison. The results show that the estimates of the effect of 3G change very little when age at survey, electricity or cultural vehicles such as TV or radio are added to the equation.

Second, a key concern in our analysis is potential measurement error in FGM data. The DHS follows a standardized protocol to maximize reliability: Women are interviewed privately about their own FGM status and that of their daughters, usually by trained female interviewers, to minimize reporting bias on this sensitive topic. However, these measures are not always perfectly implemented, and even when they are, some measurement error remains unavoidable. If misreporting of FGM is orthogonal to 3G availability, the latter would simply cause larger confidence intervals. More worryingly, if the misreporting is correlated with the availability of 3G network at the time of birth, our estimates of the effect of 3G on FGM would be overestimated. This may happen, for example, if fast internet makes women less likely to report that they have subjected their cut daughters to FGM. We believe this is not a major identification concern in our analysis. Our specification exploits within-cluster variation, comparing girls born before and after the arrival of 3G, with FGM status measured at the same time regardless of whether the girl was born before or after 3G. As a further robustness check, we re-estimate the main equation including household fixed effects, restricting variation to daughters within the same household born before or after 3G, with information for all household daughters collected at the same time from the same respondent (the mother).<sup>26</sup> The results, reported in column 4 of Table 4, remain statistically significant at conventional confidence levels, providing reassurance about our findings. Also, the following sections show that anti-FGM campaigns have no effect on support for this practice, and that the internet content driving the change is not explicitly about FGM, which is in line with the hypothesis that 3G does not change much reporting

---

<sup>26</sup>The average number of daughters per household in the analytical sample is 1.65.

behaviors.

Third, while FGM typically occurs within the first year of life in Nigeria, a non-negligible share takes place up to the third year of life (around 15% of the cuts). One may therefore wonder whether girls aged 1 or 2 at the time 3G was deployed in their community might also have had their FGM status affected by 3G. In our analysis, however, these girls were categorized as controls ( $g = -1$  and  $g = -2$ ). To address this concern, we re-estimate the results excluding girls who were 1 or 2 years old at the time of deployment. The results, reported in column 5 of Table 4 show slightly larger effects when these girls are excluded from the analytical sample.

Fourth, in Nigeria, around 80% of cuttings occur within the first year of life.<sup>27</sup> However, since we use girls aged 0–14 at the time of the survey in the analytical sample, one may wonder whether the effects could be partially capturing a delay in the age at cutting rather than fully a reduction in the prevalence. While anecdotal evidence in the field points, if anything, in the opposite direction, we test this hypothesis as follows.<sup>28</sup> We first re-estimate the main results restricting the analytical sample to girls aged at least 3 at the time of the survey. In our sample of women aged 15–49, nearly 97% of girls who will be cut were already cut by age 3. The results of this analysis are reported in Column 6 of Table 4. While the sample size reduced substantially and the point estimate is smaller, the estimated coefficient remains meaningful and statistically significant at the 5% level. Second, we estimate the effect of 3G on age at cutting. The results of this test are reported in Column 7 in Table 4. While they should be interpreted with caution because 3G might change the composition of the girls cut, the coefficient measuring the effect of 3G availability on age at cutting is small and largely insignificant.

Fifth, DHS and MICS data include the coordinates of the place of residence of survey respondents but not of the place of birth. For those households that migrate during the lifetime of the girl, the treatment variable (availability of 3G at the time of birth) might be measured with error. If this measurement error is random, it would merely bias the estimates downward, without altering the main conclusions of the analysis. The problem would however arise if fast internet affects migration (García-Hombrados et al., 2022; Adema et al., 2022), which could change the composition of villages receiving 3G. The latter may occur if the 3G network encourages the migration of more anti-FGM households into areas with fast internet, which would upwardly bias the estimated effect of interest or affect their peers. However,

---

<sup>27</sup>For example, in many Igbo communities, the cutting historically occurred during the naming ceremony on the 7th day of life.

<sup>28</sup>In a focus group in the Calabar area, an Ibibio woman highlighted that in her community, FGM is decreasing but is also being practiced earlier in life than it used to be.

this is not the case in our sample: support for FGM among migrants and non-migrants in 3G-covered areas is, if anything, slightly higher among migrants than non-migrants.<sup>29</sup> As a complementary exercise, we exploit information on migration history—available only in the 2018 DHS and 2021 MICS rounds—and reestimate the main results restricting the analytical sample to daughters whose parents have never migrated during their lifetime. The results of this analysis are reported in column 8 of Table 4 and show similar effects to those in the main analysis, suggesting that the main conclusions are not driven by a change in the composition of the individuals living in areas covered by 3G.

Sixth, one may wonder whether other policies introduced around the same time, which could potentially have affected FGM, might be confounding the effect of 3G on FGM. Such confounding could arise only if a policy coincided in timing and location with the 3G rollout, or had differential effects across areas with and without 3G. While the granular nature of the 3G expansion and the effects observed from the first year help alleviate these concerns, we explicitly test the potential confounding effect of what is arguably the policy with the largest potential impact on FGM: the staggered introduction of anti-FGM legislation in some Nigerian regions from 2015. To explore whether these laws could confound the main results, we conduct two empirical tests. First, we re-estimate the main regression controlling for whether FGM was banned in the region at the time of birth. Second, we re-estimate the main specification excluding from the sample those states where FGM was banned during the study period. The results are reported in Table A2 in Appendix A. Column 1 shows the benchmark effect from the main estimation using the full sample. Column 2 indicates that adding a dummy variable for the existence of a ban at the time of birth has virtually no effect on the point estimate. Reassuringly, column 3 shows that the coefficient remains largely unchanged when regions with FGM bans are excluded.

Seventh, the introduction of the 3G network not only provided fast internet but also facilitated communication among individuals through calls and short messages (SMS). To test whether these other smartphone functionalities could have driven the observed effects rather than internet access, we exploit the fact that 2G technology enabled calls and SMS-based communication but did not provide fast internet.<sup>30</sup> If calls or SMS were the mechanism behind the observed effects, exposure to 2G would be expected to produce similar effects on FGM. Using geolocated GSMA data on 2G coverage, we estimate the impact of the expansion of 2G mobile networks on

---

<sup>29</sup>The difference is 2 percentage points with  $p\text{-value} = 0.190$ .

<sup>30</sup>While 2G facilitated text-based communication through SMS, it only allowed limited and slow internet browsing and applications.

FGM, which occurred mainly in the early 2000s. Overall, the results reported in Table A3 and Figure A9 in Appendix A show null effects that do not increase over time. Taken together, our results suggest that the observed effects of 3G are related to the internet rather than to other communication improvements already available through 2G coverage.

Finally, a potential concern is that the observed effect may be driven not by 3G connectivity itself, but by the broader infrastructure and state presence associated with its deployment, such as access roads, electricity, or maintenance facilities. This is unlikely because 3G is typically deployed on the same towers already used for 2G, meaning that most of the physical infrastructure and accompanying state presence were already in place. Moreover, if the effects were driven by improvements in infrastructure and state presence that may accompany the construction of towers, rather than by 3G connectivity itself, we would expect to observe a similar effect from 2G coverage. As shown in the previous analysis, however, no such effect is found. To further test this hypothesis, we examine the effect of 3G availability at the time of the survey on four measures of state presence: the number of public schools opened in the county, access to electricity, and whether the respondent owns a radio or a TV. While access to electricity and ownership of radio and TV do not depend exclusively on state presence and public infrastructure, they are arguably influenced by it. We estimate equation 2 using a panel of yearly data on schools opened at the county level for the period 2007–2018, and information on access to electricity, radio, and TV from the DHS rounds 2003, 2008, 2013, and 2018. The results of these analyses are reported in Table A4 and Figure A10 in Appendix A. Although the presence of negative weights (ranging between 39% and 42%) calls for caution when interpreting the TWFE estimates, the results are consistent across methods for most measures, suggesting very limited effects of 3G availability on state presence.

## 7 Shifting the value of FGM

This section first shows that 3G reduced support for FGM and then presents evidence consistent with the hypothesis that the effect of 3G on FGM is driven by a reduction in the value of the social function served by the practice, which is the preservation of virginity and the prevention of extra-marital sex.

We start by assessing how 3G networks at the time of the survey affect support for FGM, estimating equation 2 for men aged 15–59 and women aged 15–49, the groups for which information on support for FGM is available in the DHS data. The

variable *Support for FGM* is defined as a discrete variable equal to 1 if the individual does not want the practice to continue, 2 if it depends, and 3 if the respondent wants the practice to continue. As dependent variables, we use both the discrete measure itself and a dichotomized version. The latter takes the value 0 if the individual does not want the practice to continue, and 1 if the individual answers "depends" or "yes" to the question of whether he/she wants the practice to continue.

The results of the analysis are presented in Table 5.<sup>31</sup> Since the treatment variable in this analysis is the share of population with availability of 3G network, we can only rely on the estimator developed in Chaisemartin and D’Haultfœuille (2020) and on TWFE.<sup>32</sup> While the Goodman-Bacon decomposition suggest that the Chaisemartin and D’Haultfœuille (2020) might work better in this analysis<sup>33</sup>, our findings are stable across different estimators. They reveal that 3G network availability at the time of the survey decreases the support for FGM. Our preferred estimator, reported in column (1), shows that an increase in 10 percentage points in the proportion of people covered by 3G networks decrease by 0.067 points the discrete measure of support for FGM, measured in a scale from 1 to 3, while it reduces in 3.5 percentage points the binary indicator of support for FGM. The estimated effect is sizeable, and the parallel trends tests, along with the event studies reported in Figures A11 and A12 in Appendix A, indicate that the effects are not driven by pre-existing differential trends.

The results by gender in columns 3 to 6 suggest that the observed effect of 3G on support for FGM is primarily driven by women. While the effects are large and statistically significant for women, the coefficients measuring the impact of 3G on men’s support for FGM have the opposite (positive) sign and, although sizeable, are statistically insignificant at conventional confidence levels. Nonetheless, the results for men should be interpreted with caution for two reasons. First, the male sample size is substantially smaller.<sup>34</sup> Second, Figure A11, displaying the event studies,

---

<sup>31</sup>We also present results incorporating information on support for FGM from the 2016 MICS survey. These results, reported in Table A5 in Appendix A, yield similar estimates. The main estimations reported in Table 5, however, rely exclusively on DHS data to ensure consistency in the measurement of support for FGM and in the construction of the lead and lag periods, since DHS surveys are conducted every five years. The 2021 MICS survey cannot be used, as 3G coverage data are only available up to 2018, and the treatment variable in this analysis is measured as 3G availability at the time of the survey.

<sup>32</sup>Callaway and Sant’Anna (2021); Sun and Abraham (2021) estimators can only be used with a binary treatment.

<sup>33</sup>We decompose the weights and find that 45% and 54% of the weights used in the OLS estimation are negative, which might pose challenges on the use of the TWFE model.

<sup>34</sup>Only one man in three households is interviewed in DHS rounds 2003, 2008, and 2013. Moreover, the male module in the 2018 survey round does not include information on support for FGM.

shows that men’s support for FGM seems to follow a pre-3G upward trend.

But what is driving the effect of fast internet on support for FGM and its prevalence? On the one hand, internet access might provide exposure to explicit anti-FGM content and campaigns. We test this hypothesis in detail in the following section. However, most of the online content consumed by Nigerians is unrelated to FGM. Our qualitative analysis and data from Google trends reported in Appendix C highlights that individuals in Nigeria use the internet to access cultural content such as videos, music, social media, and films. For many, access to the internet represents an unprecedented window to other cultures and women’s identities. We hypothesize that fast internet exposes these communities to cultural content and identities where premarital sex and promiscuity are less stigmatized.

We showed in Figure 3 that the main perceived benefit of FGM in Nigeria is the preservation of virginity. Consistently, the anthropological and medical literature shows that in Nigeria and other settings, cut women are perceived as more loyal to their husbands and less sexually promiscuous, traits that are highly valued in traditional communities in Nigeria (Mandara, 2001; Orubuloye et al., 2001; Gruenbaum, 2001; Mpofu et al., 2017). While not directly linked to FGM, this cultural content might change sexual behavior reducing the stigma around promiscuity and virginity, decreasing the value of the social function played by FGM and eventually eroding this harmful practice. However, while exposure to alternative values and identities might transform one’s own values to resemble more closely those of the exposed identities (La Ferrara et al., 2012; Balbo and Barban, 2014; Miho et al., 2023), it may also lead to a resurgence and strengthening of one’s own culture (Gruenbaum, 2001; Blumenstock et al., 2022).

To test this hypothesis, we first examine the extent to which fast internet affected sexual behavior. We estimate equation 2 using information reported in the DHS for men aged 15-49 and women aged 15-49 on the age at first sex, age at marriage, number of lifetime sexual partners, number of lifetime sexual partners minus number of formal unions, whether the person had an extramarital relations in the last 12 months, and the use of condom in the last sexual relation.<sup>35</sup>

Table 6 and Figures A13 and A14 in Appendix A present the results of the effect of 3G network availability at the time of the survey on these behavioral outcomes using the staggered difference-in-differences method developed in Chaisemartin and D’Haultfœuille (2020). The results for the TWFE are reported in Table A6 in Appendix A. While the TWFE estimates are less reliable due to the presence of a

---

<sup>35</sup>These were the outcomes identified in the DHS that were reported in more than a survey round that were associated with the values of purity and sexual activity only within marriage.

significant proportion of negative weights<sup>36</sup>, the main conclusions are overall consistent across estimation methods. Panel A shows the results for women. While 3G availability at the time of the survey has no effects on their age at marriage, it has a strong negative effect on the age at first sex for women, suggesting an increase in premarital sex.<sup>37</sup> Similarly, we also find that exposure to 3G network increases the reported number of lifetime sexual partners and extra-marital sex, which are also stigmatized in the traditional culture of most Nigerian communities. On the other hand, we do not observe any effect on the probability of using condom during the last sexual relation. Taken together, these results suggest that fast internet is transforming women’s sexual behavior in traditional communities towards a more progressive stance. Once again, the results of the parallel trends tests and the event studies reported in Figure A13 in Appendix A indicate that, overall, the estimates are not the result of pre-existing differential trends. Our results are also consistent with previous evidence showing that exposure to Western media culture can transform beliefs and behaviors towards more progressive views (Jensen and Oster, 2009; La Ferrara et al., 2012).

Because in many Nigerian communities sexual behaviors such as extramarital sex, premarital sex, or having multiple sexual partners are traditionally considered taboo—particularly for women—one may wonder whether the observed changes reflect greater reporting of these behaviors rather than an actual change in the behaviors themselves. While we cannot disentangle an effect on behavior from an effect on reporting, we believe that the latter could equally be an indication of cultural change, reflecting the destigmatization of these sexual behaviors. In line with this hypothesis, Table A7 in Appendix A shows that 3G also reduced the age at first sex among women older than 30, most of whom arguably had their sexual debut many years before the arrival of the internet. This result suggests that 3G affected the reporting of premarital sex, providing additional evidence in favor of the normalization of premarital sex and against the hypothesis that changes in their reported sexual behavior occurred without destigmatization.

Panel B of Table 6 reports the estimated effects of the 3G availability at the time of the survey on sexual outcomes for men. While the coefficients are sizeable, the

---

<sup>36</sup>The proportion of negative weights in the estimation of treatment effects in the TWFE for these outcomes vary between 39% and 42%, highlighting that the Chaisemartin and D’Haultfoeuille (2020) method is more appropriate in these estimations.

<sup>37</sup>An accurate measure of premarital sex cannot be constructed because age at first sex and age at marriage are reported in age-years rather than specific dates, so when marriage and first sex occur within the same age-year, it is not possible to determine whether sex occurred before marriage. However, in a context where sex after marriage is typically immediate, a reduction in the age at first sex, coupled with an increase or no change in the age at marriage, suggests an increase in premarital sex.

estimates of the effects are statistically insignificant at conventional confidence levels and the sign of the coefficients vary in each outcome. While the smaller sample of men surveyed and the results of the event study reported in Figure A14 in Appendix A require cautious interpretation, the results suggest that 3G does not make men less traditional in terms of sexual behavior. A potential explanation for the weaker cultural effects is that, as suggested by the descriptive statistics, traditional values emphasizing purity and sexual activity only within marriage were more prominent for women than for men in most Nigerian communities. Therefore, the expected shock from exposure to cultures where such behaviors are more common might be smaller for men.

While the previous analysis shows that 3G destigmatized premarital sex and shifted sexual behavior towards more liberal attitudes among women, this does not necessarily imply that the effect of fast internet on FGM operates through the destigmatization of premarital sex. To examine this mechanism more directly, we further conduct the following three analyses.

First, we estimate the effect of 3G availability on FGM in Nigerian regions where, although the practice remains significant (10% or higher), it is not primarily associated with the preservation of virginity. In these areas, the most cited benefits of FGM include social acceptance (e.g., rite of passage) and false beliefs about hygiene or religion, while the preservation of virginity is mentioned only sparsely as a benefit of FGM (less than 3%). We identify the relevant placebo regions using information on FGM motives included in the 2008 DHS survey, years before the arrival of 3G to Nigeria. If the main mechanism through which fast internet affects FGM is the destigmatization of premarital sex, we should not observe an effect in areas where FGM is not linked to premarital sex. The results of this placebo analysis are reported in Column (6) Table 3. Reassuringly, they show that fast internet has no effect on FGM in regions where FGM is unrelated to the preservation of virginity.

Second, in a similar spirit, we explore whether the effect of 3G availability is stronger, the more linked FGM is to the preservation of virginity. To conduct this analysis, we expand equation 1 and add an interaction of the 3G availability variable with a regional indicator of the degree to which FGM is linked to the preservation of virginity in 2008.<sup>38</sup> In line with this mechanism, the results reported in column 4 of Table 3 show that the interaction term is negative and largely significant, showing that the effect of fast internet on FGM is particularly large in areas in which this practice is more strongly linked to the preservation of virginity.

---

<sup>38</sup>The regional indicator is the share of individuals in the region that declare a benefit of FGM is that it preserves virginity.

Finally, we conduct a placebo analysis of the effect of 3G availability in Nigerian regions where out-of-marriage sex was common in 2008, before the arrival of fast internet in the country.<sup>39</sup> If out-of-marriage sex was already normalized, the effect of the internet on FGM is unlikely to be driven by a destigmatization of premarital sex in those areas.

Using data from the 2008 DHS survey, we calculate the intensity of out-of-marriage relations at the regional level by computing, for each woman in the women’s module, the number of lifetime sexual partners minus the number of marriage unions, and averaging the result across states. The average varies substantially, ranging from 0.02 to 4.8. We define states where premarital sex is normalized as those where the average number of out-of-marriage relations is 2 or higher. These include Abia, Akwa Ibom, Benue, Delta, Ekiti, Enugu, Imo, and Rivers. While the prevalence of FGM is negatively correlated with the intensity of out-of-marriage relations, the practice is still existing in many of these states, reassuring that, at least in some areas, avoiding premarital sex is likely not the only reason for FGM. The results of this analysis are reported in column 7 of Table 3 and show that the coefficient measuring the effect of 3G availability on FGM in these regions is very small and statistically insignificant. The results do not change substantially when we use different thresholds to define regions with normalized out-of-marriage sex.<sup>40</sup> These results confirm that, in those areas in which premarital sex was already destigmatized before the arrival of 3G, 3G had little effect.

All in all, we find evidence not only that fast internet transformed sexual behavior toward more liberal attitudes, but also that the diminishing influence of traditional values associated with FGM—such as women’s virginity and sexual loyalty to their husbands—is driving the effect of fast internet on FGM.

## 8 Alternative hypotheses

### 8.1 The effect of explicit anti-FGM content on-line

The effects of fast internet on the support for FGM and on the prevalence of FGM might also be driven by exposure to explicit anti-FGM online content, rather than

---

<sup>39</sup>Another obvious placebo analysis would rely on information from the Murdock Atlas regarding ethnic groups in which premarital sex has not traditionally been stigmatized. In the case of Nigeria, however, only two minor groups meet this criterion: the Wodaabe and the Maguzawa. This analysis cannot be conducted because these ethnicities are not identified in the DHS or MICS surveys used in the analysis.

<sup>40</sup>The effect using a threshold of 1.5 is nearly identical. While the effect at a threshold of 1 is larger, it remains much smaller than in regions where the number is below 1.

by reducing premarital sex stigma. Indeed, the internet facilitates the dissemination of explicit content related to FGM, granting individuals the opportunity to access such materials. Nigeria stands out globally for its substantial efforts in countering FGM through online platforms and social media channels.<sup>41</sup> Approximately 14% of all FGM-related tweets originated from Nigerian users, and while not all online discussions on FGM are explicitly critical of the practice, the vast majority are (Babbs et al., 2023).

To better understand what effect explicit anti-FGM online content and campaigns could have, we assess the short-term effects on support for FGM of the anti-FGM online campaign for the International Day of the Girl Child, celebrated on October 11, 2018.<sup>42</sup> This campaign is launched on the same date every year by UNICEF in collaboration with the Nigerian Ministries of Health and Social Welfare; and of Women Affairs and Social Development. Through these collective efforts, the campaign strives to engage a broad audience. Figures A16 and A17 in Appendix A show that October 11th is the day of the year with a higher number of anti-FGM tweets and that Google searches about FGM peak every year on October 11th.

We focus in the 2018 edition of the campaign because it was launched in the middle of the 2018 round of the Nigerian DHS survey fieldwork, which lasted from August to December 2018.<sup>43</sup> To estimate the short term effects of the campaign, we use a regression discontinuity design (RDD) in time, which compares the support for FGM among those women aged 15-49 interviewed just before and after October 11th 2018 in areas where 3G is available. Because the male module of the 2018 survey round does not include questions on support for FGM, the analysis is conducted only for women. Specifically, we estimate the following regression:

$$Supports\ FGM_i = \alpha_0 + \alpha_1 Post_i + \alpha_2 F(Days\ to\ October\ 11th_i) + \mu_i \quad (3)$$

where *Supports FGM* is the main outcome and indicates the degree of FGM support.

---

<sup>41</sup>For example, the “Sarah’s Cross” series, available on YouTube, successfully reached an audience exceeding 200,000 individuals within Nigeria. The #endcuttinggirls Twitter campaign, spanning various social media platforms, effectively engaged nearly 8 million people across the country. Additionally, an anti-FGM documentary titled “Female Genital Mutilation in Nigeria” garnered 600,000 views (UNFPA-UNICEF, 2018). In the year 2020, Nigeria demonstrated heightened online activism against FGM, emerging as the second country, following Kenya, to generate a significant volume of tweets discussing the issue.

<sup>42</sup>Examples of anti-FGM tweets issued the October 11, 2018 are provided in Figure A15 in Appendix A. Key hashtags like #endcuttinggirls or #EndFGM trend on Twitter and Instagram, amplified the campaign’s reach.

<sup>43</sup>The previous DHS survey was launched in 2013, before the start of the anti-FGM online campaigns for the International Day of the Girl’s Child in Nigeria.

The variable *Post* is a dummy variable equal to 1 if woman  $i$  is interviewed on the 11th of October 2018 or later and 0 if is interviewed before the mentioned date. The main parameter of interest is  $\alpha_1$ , which measures the effect of being interviewed just after the anti-FGM online campaign on October 11th.  $F(\text{Days to October 11th})$  is a function of the number of days between the survey interview and October 11th, and  $\mu$  is the error term. Equation 3 is estimated using both non-parametric methods and parametric methods with first and second order polynomials for the running variable. The non-parametric estimation is calculated using the local polynomial regression-discontinuity estimation method with robust bias-corrected confidence intervals developed in Calonico et al. (2019). The parametric regressions include survey cluster fixed effects. As a robustness check, we also estimate these regressions excluding from the analysis those women that were interviewed during the week before the start of the campaign to examine the robustness of the results to potential anticipation effects during pre-campaign days.

The results of this analysis are presented in Table 7. They show null short-term effects of the campaign on the support for FGM for individuals living in areas with 3G network available. The estimates are consistently small and statistically insignificant at conventional confidence levels across estimation methods and samples. Figure 7 shows graphically the effect of the October 11th campaign for individuals with 3G network available in their location, showing small and largely insignificant coefficients.<sup>44</sup> The graph also shows null effects for those women that live in areas with no 3G network available, which serves as a placebo test. While the RDD approach used only measures the short-term effects of the anti-FGM campaign, it is likely that if these campaigns have no impact in the short term, the long-term effects would also be limited.

## 8.2 Income shock

Fast internet may reduce the prevalence of FGM even if 3G availability does not diminish support for the practice. If poverty plays a pivotal role in FGM, and parents cut their daughters to secure a higher bride price (Khalifa, 2022; García-Hombrados and Salgado, 2022), then a positive effect of fast internet on income may lead some parents at the margin to refrain from cutting their daughters, even if their support for FGM does not change. This mechanism is unlikely in Nigeria, where FGM is not typically discussed in bride price negotiations and where communities with near-universal prevalence of FGM are rare. Our cumulative suggestive evidence

---

<sup>44</sup>Figure A18 in Appendix A shows similar effects for the dichotomous measure of support for FGM.

indicates that, if anything, this factor plays only a limited role.

To assess this mechanism empirically, we first examine the economic effects of 3G using DHS data. The results, reported in columns 1 and 2 of Table A8 in Appendix A, suggest that 3G coverage did not increase household wealth.<sup>45</sup> Our results, however, contrast with those of Bahia et al. (2024), who, using a different dataset, find that 3G coverage increases consumption and labor force participation, and reduces poverty in Nigeria.

Thus, while our results show no effect of 3G coverage on wealth, we cannot rule out a positive economic effect, given previous evidence and the fact that the DHS may not be the most suitable dataset to measure key economic dimensions such as income or consumption. We therefore propose a different strategy to test this mechanism: we examine the effect of rainfall shocks, which crucially affect economic welfare. If the effect of 3G on FGM were purely driven by income shocks caused by 3G, other shocks affecting economic welfare—such as rainfall shocks—should similarly affect the prevalence of FGM. Specifically, we replicate the multi-country analysis conducted by McGavock and Novak (2023) focusing on Nigeria. The latter study examines the effect of adverse rainfall shocks during the age of highest risk of cutting on the prevalence of FGM. The results, reported in Table D1 in Appendix D, show that negative income shocks do not increase the prevalence of FGM in Nigeria. This finding is consistent with McGavock and Novak (2023), who report no effect of rainfall shocks on FGM among ethnic groups where the practice is typically conducted early in life, as in the case of Nigeria. A detailed description of this analysis is provided in Appendix D. Overall, these results suggest that, if existent, potential improvements in income or wealth caused by 3G are unlikely to drive the observed effect of 3G on FGM.

### 8.3 Decreased religiosity

One may wonder whether the reduction in FGM is driven by 3G availability decreasing religiosity. Although the practice of FGM is not theologically grounded in any major religious group (Adewole and Adayonfo, 2017; Burrage, 2016), some communities across Africa perceive certain forms of FGM as being religiously motivated.<sup>46</sup> In Nigeria, DHS data from 2018 show that 10% of women aged 15–49 believe that

---

<sup>45</sup>The results of the parallel trends tests reported in Table A8 in Appendix A, together with the event studies shown in Figures A19 and A20, suggest that the findings are not driven by pre-existing differential trends. While 3G does not appear to increase women’s labor force participation, Table A9 in Appendix A shows that 3G increases bargaining power for women and reduces it for men.

<sup>46</sup>This is the case, for example, with the Sunna mutilation practiced among Somali groups in East Africa (Mehari, 2023).

FGM is a practice linked to religion.

While the proportion of individuals in Nigeria who report having no religion is very small (below 1%)<sup>47</sup>, one might argue that fast internet reduces the intensity of religiosity, potentially weakening the practice of FGM if it is motivated mainly by religious reasons. To test this hypothesis, we use individual-level DHS data for women aged 15–49, which includes a question on whether they believe that FGM is a religious requirement. We use this information to estimate the differential effect of 3G on FGM by whether the girl’s mother believes that FGM is a religious requirement. If the reduction in FGM were driven by decreasing religiosity, we should observe a significant decline in the prevalence of FGM precisely among those who believe it is religiously mandated.

The results of this analysis are reported in column 5 of Table 3. They show that the interaction term *3G availability*  $\times$  *FGM linked to religion* is positive and has the same magnitude but the opposite sign as the main *3G availability* coefficient. While the interaction is not statistically significant, it suggests that the overall effect is driven by households that do not believe FGM is a religious requirement, whereas the effect among those who do believe so is small and largely statistically insignificant. Taken together, these results suggest that the effects of 3G on FGM are unlikely to be driven by a decline in religiosity.

## 8.4 Increased in education

Finally, one may wonder whether internet was used for educational purposes, potentially increasing educational attainment. While there is a negative correlation between maternal education and FGM, existing causal evidence from Nigeria shows that the expansion of women’s education has no effects on their support for FGM and on the prevalence of FGM among their daughters (De Cao and La Mattina, 2019). To explore this potential mechanism in more detail, we assess the effect of 3G on education. We conduct two different analyses.

First, focusing on the sample of individuals aged 18 or older, we estimate the effect of 3G on educational attainment and on taking secondary education by measuring the number of years of exposure to 3G during the educational period, which in Nigeria lasts from age 6 (when students start compulsory education) to age 18 (when individuals who do not repeat grades complete secondary education). Specifically, we estimate the following regression:

---

<sup>47</sup>This information comes from the sample of men and women surveyed in the 2018 round of the DHS.

$$SchoolAttainment_{ibv} = \delta_0 + \delta_1 Years\ Exposed\ to\ 3G_{bv} + \theta b + \gamma_v + u_{ibv} \quad (4)$$

where  $SchoolAttainment_{ibv}$  is the educational attainment of individual  $i$ , aged  $b$  and living in village  $v$ . The  $SchoolAttainment$  variable takes the value 0 if the individual has no education, 1 if primary education, 2 if secondary education, and 3 if higher education.  $Years\ Exposed\ to\ 3G$  indicates the number of years the individual was exposed to 3G during his/her educational years, which in Nigeria go from age 6 to age 18. In principle, the variable can range between 0 and 13 years and varies at the village  $v$  and year-of-birth  $b$  level.  $\delta_1$  is the parameter of main interest and measures the effect of one additional year of 3G network availability during the educational years on educational attainment or on taking secondary education.  $\theta_b$  and  $\gamma_v$  are year-of-birth and village fixed effects to control for time-invariant village-level and year-of-birth specific events. Standard errors are clustered at the village level (DHS survey cluster). We also estimate the regression using a binary dependent variable indicating whether the individual has taken secondary education.

The results of this analysis are reported in columns 1 and 2 of Table A10 in Appendix A.<sup>48</sup> The coefficients are negative and small in magnitude, but statistically significant at conventional confidence levels, suggesting that, if anything, 3G reduces educational attainment and secondary education rather than increasing it.

As a second exercise to test the effect of 3G availability on educational outcomes, we also estimate equation 2 using school enrollment at the time of the survey as the dependent variable, using information available on school enrollment for individuals aged 6–18 at the time of the survey.<sup>49</sup> The regressions are estimated using TWFE and the estimation method for staggered difference-in-differences with a continuous treatment variable developed in Chaisemartin and D’Haultfoeuille (2020) for the full sample, girls, and boys. The results of this exercise are reported in columns 3 and 4 of Table A10. The coefficients are small and, overall, largely statistically insignificant for both men and women.

Overall, these analyses suggest that 3G availability did not improve educational outcomes, making it unlikely that it reduced FGM via increases in maternal or paternal education.

---

<sup>48</sup>The event study for the effect of 3G network coverage on school attendance is reported in Figure A21. The figure confirms that treatment and control observations followed parallel trends prior to the deployment of 3G networks.

<sup>49</sup>In DHS surveys, information on school enrollment is provided in the household roster for all individuals in the household regardless of their age.

## 9 What internet content is driving the change in sexual behavior?

Existing databases in Nigeria lack granular information on the specific internet content accessed, which prevents a comprehensive quantitative analysis of which particular content is driving the effects of 3G on FGM and sexual behavior. To explore this issue and gain a better understanding of cultural change and its drivers in Nigeria, we implemented the following analyses.

First, we conducted a series of focus group discussions in 8 rural communities across Cross Rivers and Abuja states in Nigeria, including 68 participants from the Efok, Ekoi, Ibibio, Hausa, Yoruba, and Igbo ethnic groups. The goal of these discussions was to understand (1) how individuals used the internet and what kind of content they accessed; (2) the evolution of norms around sex in the community; and (3) the perceived effects of internet on social norms, particularly on sexual behavior and stigmas. Here we provide a brief summary of the main results. Further details about the design and the results of the qualitative analysis are reported in Appendix C.

Across groups, responses were remarkably consistent: internet was primarily used for communication, entertainment (especially movies, music, and—particularly among younger participants—social media), and accessing news. Participants across all communities identified both positive and negative transformations in their communities associated with internet access. On the positive side, improvements in access to entertainment, communication with family members living outside the community, and economic opportunities were frequently mentioned. On the negative side, participants expressed concerns about online fraud and moral decline. Foreign movies, social media, and pornography were seen as major drivers of cultural erosion. A common theme across all groups was that young people were increasingly imitating behaviors seen online—becoming more selfish, more prone to violence, and engaging in sexual relationships before marriage.

Regarding premarital sex, participants unanimously agreed that significant changes had occurred in recent years. What was once socially unacceptable has become widely normalized, especially among youth. In southern communities, traditional leaders appeared to have largely adapted to the shift, whereas in northern communities the issue is provoking notable intergenerational tension, with elders voicing strong concern. Across all focus groups, content depicting premarital sex as normal, as well as content explicitly showing naked bodies and sexual activity, was cited as the primary cause of the shift in sexual norms, with no dissenting voices on this

point.

Second, we provide data from Google Trends on Google searches in Nigeria. Our results indicate that football, sports betting, social media, YouTube, websites that provide free access to films, as well as “sex” and various “pornographic” terms and websites, rank among the country’s top search terms in Google, with several million searches each month.

Taken together, the results from Google Trends and the focus groups suggest that Nigerians regularly access content depicting premarital sex as normal and content displaying explicit sexual images. This exposure is perceived as driving changes in sexual behaviors and reducing the stigma surrounding premarital sex. While we cannot empirically test whether these contents are indeed triggering the observed changes, previous evidence showed that cultural content portraying Western lifestyles can shift beliefs and behaviors toward more progressive views (Jensen and Oster, 2009; La Ferrara et al., 2012), and that exposure to sexually explicit content can change sexual behaviors (Brown et al., 2006; Peter and Valkenburg, 2008).

While access to sexually explicit content is perceived by the local population as a key contributor to the normalization of premarital sex, with potential implications for FGM, we are not making normative judgments about the desirability of promoting such content, as this could not only raise ethical concerns but also have unintended negative effects on mental health and aggressive behavior (Waterman et al., 2022; Wright et al., 2016; Zhou et al., 2021; Setyawati et al., 2020). Nonetheless, the results of this study are also valuable for policy: Beyond this specific case, they suggest that interventions aiming to challenge harmful norms should focus on reducing associated stigmas in order to diminish the social value of these norms.

## 10 Conclusions

Harmful social norms such as FGM, breast ironing, or child marriage continue to affect millions of women worldwide. FGM alone threatens nearly four million girls each year and while the prevalence of these practices is declining in some areas, eradication is far from imminent.

What can be done to accelerate this process? Policy responses have often relied on advocacy campaigns that stress the health risks of these practices or on education programs. Yet, evidence shows that these approaches have limited effects—suggesting also that parents are not necessarily unaware of the harms involved. Instead, these norms persist because they fulfill important social functions.

In the case of Nigeria, line with anthropological research, our study provides de-

scriptive evidence that, in Nigeria, FGM is believed to discourage women from engaging in premarital or extramarital relations, a function that becomes particularly salient in contexts where such practices are highly stigmatized. We then examine the impact of one of the most significant cultural shocks in recent decades in sub-Saharan Africa: the expansion of fast internet. This technological shift brought new cultural content and alternative models of women's behavior to traditional communities. We find that exposure to 3G networks leads to both a decline in the prevalence of FGM and diminished support for the practice. Our results also suggest that this reduction in FGM is driven by 3G fostering more progressive sexual norms and reducing premarital sex stigma, thereby diminishing the social value of FGM as a mechanism of controlling sexuality.

While fast internet may also entail adverse effects, such as cultural convergence and other unintended outcomes, our findings highlight that harmful practices can decline when the social value of the function they serve is eroded. Beyond 3G, our results suggest that interventions targeting the stigmas and beliefs sustaining these norms may be more effective than efforts aimed directly at the harmful practices themselves.

## References

- Adema, J. A. H., Aksoy, C. G., and Poutvaara, P. (2022). Mobile Internet Access and the Desire to Emigrate. CESifo Working Paper 9758.
- Ademiluka, S. O. (2023). And when i came to her i found she was not a virgin. *Verbum et Ecclesia*, 44(1):1–10.
- Adewole, A. J. and Adayonfo, E. O. (2017). Correlates of female genital mutilation (fgm) among patients of a teaching hospital in Nigeria. *IFE Psychologia: An International Journal*, 25(1):223–231.
- Ahmadu, F. (2001). Rites and wrongs: An insider/outsider reflects on power and excision. In Shell-Duncan, B. and Hernlund, Y., editors, *Female Circumcision in Africa: Culture, Controversy and Change*, pages 283–312. Lynne Rienner Publishers.
- Ahmed, R., Brimble, P., Kovvuri, A., Saia, A., and Yang, D. (2025). Ancient Epics in the Television Age: Mass Media, Identity, and the Rise of Hindu Nationalism in India. Working Paper Series 33417, National Bureau of Economic Research.
- Babbs, G., Weber, S., Abdalla, S., Cesare, N., and Nsoesie, E. (2023). Use of machine learning methods to understand discussions of female genital mutilation/cutting on social media. *PLOS global public health*, 3.
- Bahia, K., Castells, P., Cruz, G., Masaki, T., Pedros, X., Pfutze, T., Rodriguez-Castelan, C., and Winkler, H. (2024). The welfare effects of mobile broadband internet: Evidence from Nigeria. *Journal of Development Economics*, 170.
- Balbo, N. and Barban, N. (2014). Does fertility behavior spread among friends? *American Sociological Review*, 79(3):412–431.
- Becker, A. (2024). On the economic origins of concerns over women’s chastity. *The Review of Economic Studies*, 92(4):2303–2329.
- Bellemare, M., Novak, L., and Steinmetz, T. L. (2015). All in the family: Explaining the persistence of female genital cutting in West Africa. *Journal of Development Economics*, 116(C):252–265.
- Berg, R., Underland, V., Odgaard-Jensen, J., Fretheim, A., and Vist, G. (2014). Effects of female genital cutting on physical health outcomes: A systematic review and meta-analysis. *BMJ open*, 4.

- Bicchieri, C. (2005). *The Grammar of Society: The Nature and Dynamics of Social Norms*. Cambridge University Press.
- Blumenstock, J., Dube, O., and Hussain, K. (2022). Can secular media create religious backlash? evidence from pakistan’s media liberalization. EDI Working Paper Series.
- Brown, J., L’Engle, K., Pardun, C., Guo, G., Kenneavy, K., and Jackson, C. (2006). Sexy media matter: Exposure to sexual content in music, movies, television, and magazines predicts black and white adolescents’ sexual behavior. *Pediatrics*, 117(4):1018–1027.
- Burrage, H. (2016). *Female Mutilation: The Truth Behind the Horrifying Global Practice of Female Genital Mutilation*. New Holland Publishers.
- Callaway, B. and Sant’Anna, P. H. (2021). Difference-in-differences with multiple time periods. *Journal of Econometrics*, 225(2):200–230.
- Calonico, S., Cattaneo, M. D., Farrell, M. H., and Titiunik, R. (2019). Regression Discontinuity Designs Using Covariates. *The Review of Economics and Statistics*, 101(3):442–451.
- Camilotti, G. (2015). Interventions to Stop Female Genital Cutting and the Evolution of the Custom: Evidence on Age at Cutting in Senegal. *Journal of African Economies*, 25(1):133–158.
- Cetorelli, V., Wilson, B., Batyra, E., and Coast, E. (2020). Female genital mutilation/cutting in Mali and Mauritania: Understanding trends and evaluating policies. *Studies in Family Planning*, 51(1):51–69.
- Chaisemartin, C. and D’Haultfoeuille, X. (2020). Two-way fixed effects estimators with heterogeneous treatment effects. *American Economic Review*, 110(9):2964–2996.
- Chong, A. and Ferrara, E. L. (2009). Television and Divorce: Evidence from Brazilian Novelas. *Journal of the European Economic Association*, 7(2/3):458–468.
- Congdon Fors, H., Isaksson, A.-S., and Lindskog, A. (2024). Changing local customs: The long run impacts of Christian missions on female genital cutting in Africa. *Journal of Development Economics*, 166.
- Corno, L. and La Ferrara, E. (2019). Changing harmful norms: Evidence from a randomized controlled trial on female genital cutting. AEA RCT Registry.

- Crisman, B., Dykstra, S., Kenny, C., and O'Donnell, M. (2016). The Impact of Legislation on the Hazard of Female Genital Mutilation/Cutting: Regression Discontinuity Evidence from Burkina Faso. Center for Global Development Working Paper 432.
- De Cao, E. and La Mattina, G. (2019). Does maternal education decrease female genital cutting? *AEA Papers and Proceedings*, 109:100–104.
- Diabate, I. and Mesplé-Somps, S. (2019). Female genital mutilation and migration in Mali: Do return migrants transfer social norms? *Journal of Population Economics*, 32(4):1125–1170.
- Efferson, C., Vogt, S., Elhadi, A., Ahmed, H. E. F., and Fehr, E. (2015). Female genital cutting is not a social coordination norm. *Science*, 349:1446–1447.
- Fernández, R. (2025). Understanding cultural change. NBER Working Papers 34077, National Bureau of Economic Research.
- Frezza, E. (2023). Fast Internet and Gender Norms: Evidence from Sub-Saharan Africa. Working paper.
- García-Hombrados, J., Ciacci, R., and Zainudeen, A. (2022). Mobile phone network and migration: Evidence from myanmar. *International Migration Review*, 56(2):594–614.
- García-Hombrados, J. and Salgado, E. (2022). Legal Bans, Female Genital Cutting, and Education: Evidence from Senegal. *The World Bank Economic Review*, 37(1):74–92.
- Giuliano, P. and Nunn, N. (2020). Understanding cultural persistence and change. *The Review of Economic Studies*, 88(4):1541–1581.
- Goodman-Bacon, A. (2021). Difference-in-differences with variation in treatment timing. *Journal of Econometrics*, 225(2):254–277.
- Gruenbaum, E. (2001). *The female circumcision controversy: an anthropological perspective*. University of Pennsylvania Press.
- Gulesci, S., Jindani, S., La Ferrara, E., Smerdon, D., Sulaiman, M., and Young, H. (2023). A stepping stone approach to norm transitions. HKS Working Paper RWP23-013.

- Gulesci, S., La Ferrara, E., Smerdon, D., and Sulaiman, M. (2020). Changing harmful norms through information and coordination: Experimental evidence from somalia. AEA RCT Registry.
- Guriev, S., Melnikov, N., and Zhuravskaya, E. (2020). 3G Internet and Confidence in Government. *The Quarterly Journal of Economics*, 136(4):2533–2613.
- Harari, M. (2019). Women’s Inheritance Rights and Bargaining Power: Evidence from Kenya. *Economic Development and Cultural Change*, 68(1):189–238.
- Henrich, J. (2015). *The Secret of Our Success: How Culture Is Driving Human Evolution, Domesticating Our Species, and Making Us Smarter*. Princeton University Press.
- Henrich, J. (2020). *The WEIRDest People in the World: How the West Became Psychologically Peculiar and Particularly Prosperous*. Farrar, Straus and Giroux.
- Henrich, J., Boyd, R., Bowles, S., Camerer, C., Fehr, E., Gintis, H., and McElreath, R. (2001). In search of homo economicus: Behavioral experiments in 15 small-scale societies. *American Economic Review*, 91(2):73–78.
- Henrich, J. and Muthukrishna, M. (2021). The origins and psychology of human cooperation. *Annual Review of Psychology*, 72:207–240.
- Hjort, J. and Poulsen, J. (2019). The Arrival of Fast Internet and Employment in Africa. *American Economic Review*, 109(3):1032–79.
- Jensen, R. and Oster, E. (2009). The Power of TV: Cable Television and Women’s Status in India\*. *The Quarterly Journal of Economics*, 124(3):1057–1094.
- Jones, H., Diop, N., Askew, I., and Kaboré, I. (1999). Female genital cutting practices in Burkina Faso and Mali and their negative health outcomes. *Studies in Family Planning*, 30(3):219–230.
- Katz, I., Sanders, R., Carvalho, M. N., Friedman, H. S., Legesse, B., Winfrey, W., and Diop, N. (2021). Cost and impact of scaling up female genital mutilation prevention and care programs: estimated resource requirements and impact on incidence and prevalence. *PLOS ONE*, 16(1).
- Khalifa, S. (2022). Female genital cutting and bride price. *Working Paper*.
- Koso-Thomas, O. (1987). *The Circumcision of Women: A Strategy for Eradication*. Zed Books.

- La Ferrara, E., Chong, A., and Duryea, S. (2012). Soap operas and fertility: Evidence from Brazil. *American Economic Journal: Applied Economics*, 4(4):1–31.
- La Ferrara, E., Corno, L., and Voena, A. (2020). Female genital cutting and the slave trade. CEPR Discussion Papers 15577.
- Manacorda, M. and Tesei, A. (2020). Liberation Technology: Mobile Phones and Political Mobilization in Africa. *Econometrica*, 88(2):533–567.
- Mandara, M. (2001). Female genital cutting in Nigeria: Views of Nigerian doctors on the medicalization debate. In Shell-Duncan, B. and Hernlund, Y., editors, *Female Circumcision in Africa: Culture, Controversy and Change*. Lynne Rienner Publishers.
- McGavock, T. and Novak, L. (2023). Now, Later, or Never? Evidence of the effect of weather shocks on female genital cutting in Sub-Saharan Africa. *Journal of Development Economics*, 165.
- Mehari, G. (2023). Ending female genital mutilation: Progress and challenges in the Somali region, Ethiopia. In Nakamura, K., Miyachi, K., Miyawaki, Y., and Toda, M., editors, *Female Genital Mutilation/Cutting: Global Zero Tolerance Policy and Diverse Responses from African and Asian Local Communities*, pages 57–72. Springer Nature Singapore.
- Miho, A., Jarotschkin, A., and Zhuravskaya, E. (2023). Diffusion of Gender Norms: Evidence from Stalin’s Ethnic Deportations. *Journal of the European Economic Association*, 22(2):475–527.
- Mpofu, S., Odimegwu, C., Wet, N. D., Adedini, S., and Akinyemi, J. (2017). The relation of female circumcision to sexual behavior in Kenya and Nigeria. *Women & Health*, 57(7):757–774.
- Novak, L. (2020). Persistent norms and tipping points: The case of female genital cutting. *Journal of Economic Behavior and Organization*, 177:433–474.
- Novak, L. and Bussberg, C. (2023). The complexities of female genital cutting: Understanding maternal influence, stated benefits, and policy design. Working paper.
- Orubuloye, I., Caldwell, P., and Caldwell, J. (2001). Female circumcision among the Yoruba of southwest Nigeria: the beginning of change. In Shell-Duncan, B. and Hernlund, Y., editors, *Female Circumcision in Africa: Culture, Controversy and Change*. Lynne Rienner Publishers.

- Pathmendra, P., Raggatt, M., Lim, M., Marino, J., and Skinner, R. (2023). Exposure to pornography and adolescent sexual behavior: Systematic review. *Journal of Medical Internet Research*, 25.
- Peter, J. and Valkenburg, P. (2008). Adolescents' exposure to sexually explicit internet material and sexual preoccupation: A three-wave panel study. *Media Psychology*, 11.
- Poyker, M. (2023). Regime stability and the persistence of traditional practices. *The Review of Economics and Statistics*, 105(5):1175–1190.
- Setyawati, R., Hartini, N., and Suryanto, S. (2020). The psychological impacts of internet pornography addiction on adolescents. *Humaniora*, 11:235–244.
- Shell-Duncan, B. and Hernlund, Y. (2001). *Female Circumcision in Africa: Culture, Controversy and Change*. Lynne Rienner Publishers.
- Sun, L. and Abraham, S. (2021). Estimating dynamic treatment effects in event studies with heterogeneous treatment effects. *Journal of Econometrics*, 225(2):175–199.
- Taiwo, M. O., Oyekenu, O., and Hussaini, R. (2023). Understanding how social norms influence access to and utilization of adolescent sexual and reproductive health services in northern nigeria. *Frontiers in Sociology*, 8.
- UNFPA-UNICEF (2018). Performance Analysis for Phase II. UNFPA-UNICEF Joint Programme on Female Genital Mutilation.
- UNICEF (2024). Female Genital Mutilation: A Global Concern. Available from: <https://data.unicef.org/resources/female-genital-mutilation-a-global-concern-2024/>.
- USAID (2011). Female Genital Cutting Module: Questionnaire and Interviewer's Manual. Available from: [https://dhsprogram.com/pubs/pdf/DHSQMP/DHS6\\_Module\\_Female\\_Genital\\_Cutting\\_19Aug2013\\_DHSQMP.pdf](https://dhsprogram.com/pubs/pdf/DHSQMP/DHS6_Module_Female_Genital_Cutting_19Aug2013_DHSQMP.pdf).
- Vandenbosch, L. and Eggermont, S. (2015). Sexualization of adolescent boys: Media exposure and boys' internalization of appearance ideals, self-objectification and body surveillance. *Men and Masculinities*, 18(4):483–507.
- Vogt, S., Ahmed Mohammed Zaid, N., El Fadil Ahmed, H., Fehr, E., and Efferson, C. (2016). Changing cultural attitudes towards female genital cutting. *Nature*, 538.

- Wagner, N. (2015). Female Genital Cutting and Long-Term Health Consequences - Nationally Representative Estimates across 13 Countries. *Journal of Development Studies*, 51(3):226–246.
- Waterman, E., Wesche, R., Morris, G., Edwards, K., and Banyard, V. (2022). Prospective associations between pornography viewing and sexual aggression among adolescents. *Journal of Research on Adolescence*, 32.
- WHO (2025). Female Genital Mutilation. WHO Fact Sheets. Available from: <https://www.who.int/news-room/fact-sheets/detail/female-genital-mutilation>.
- World Bank (2024). Fixed broadband subscriptions (per 100 people) - Nigeria. data retrieved from World Development Indicators, <https://data.worldbank.org/indicator/IT.NET.BBND.P2?locations=NG>.
- Wright, P. J., Tokunaga, R. S., and Kraus, A. (2016). A meta-analysis of pornography consumption and actual acts of sexual aggression in general population studies. *Journal of Communication*, 66(1).
- Zhou, Y., Liu, T., Yan, H. Y., and Paul, B. (2021). Pornography use, two forms of dehumanization, and sexual aggression: Attitudes vs. behaviors. *Journal of Sex & Marital Therapy*, 47.
- Zhuravskaya, E., Petrova, M., and Enikolopov, R. (2020). Political effects of the internet and social media. *Annual Review of Economics*, 12(1):415–438.

Table 1: Descriptive statistics

PANEL A: Sample of females used in the FGM analysis					
	N	Mean	SD	Min	Max
FGM (0/1)	64,018	0.169		0	1
3G at time of birth (0/1)	64,018	0.105		0	1
Age at survey	64,018	4.900	3.493	0	14
Year of birth	64,018	2011.815	3.177	2007	2018
PANEL B: Sample of men and women used in the analysis of support for FGM, sexual behavior, and other outcomes					
	(1)	(2)	(3)	(4)	(5)
	N	Mean	SD	Mean men	Mean women
<i>Support for FGM analysis</i>					
Support for FGM (1-3)	83,043	1.542	0.847	1.564	1.534
1.No (0/1)	83,043	0.693		0.69	0.694
2. Depends (0/1)	83,043	0.072		0.057	0.078
3. Yes (0/1)	83,043	0.235		0.254	0.228
Support for FGM (0/1)	83,043	0.307		0.310	0.306
Age	83,043	30.757	9.936	32.753	30.037
Female (0/1)	83,043	0.735			
Share pop with 3G available	83,043	0.090	0.231	0.037	0.110
<i>Sexual behavior, LFP and wealth analyses</i>					
Age at 1st marriage	117,792	19.634	5.944	25.287	17.938
Age at 1st intercourse	131,673	17.744	4.069	20.324	16.822
Lifetime sexual partners	156,611	1.961	4.681	3.174	1.480
Lifetime sexual partners (excluding spouses)	153,740	1.105	4.583	2.409	0.621
Sexual partners excluding spouse (last 12 m.)	168,531	0.147	0.524	0.256	0.104
Used condom last relation (0/1)	124,647	0.094		0.177	0.064
Labor force partic. (0/1)	168,568	0.672		0.811	0.616
Hh wealth index	169,172	0	1	0.044	-0.018
Female (0/1)	169,172	0.715			
Age	169,172	29.458	10.186	30.984	28.851
Share pop with 3G available	169,172	0.115	0.255	0.107	0.119
<i>Internet access analysis</i>					
Internet access (0/1)	52,759	0.154		0.196	0.116
Share pop with 3G available	54,335	0.184	0.387	0.187	0.181

*Note:* Panel A reports descriptive statistics for the main analytical sample used to study the effect of 3G availability at birth on FGM. The sample consists of girls born between 2007 and 2018 (years for which 3G data are available), whose mothers were interviewed in the 2013 and 2018 DHS rounds and the 2016–17 and 2021 MICS rounds. FGM status for these daughters is reported by their mothers as they are younger than 15 at the time of the survey. Panel B reports descriptive statistics for the analysis of 3G at the time of the survey on support for FGM, sexual behavior, and other outcomes. This sample includes women aged 15–49 and men aged 15–59 interviewed in the 2003, 2008, 2013, and 2018 DHS rounds. Differences in the number of observations across variables, and between this table and the estimation tables, are due to missing values, the fact that the FGM module is not applied in all surveyed households, the lack of information on some outcomes in certain DHS rounds, and the exclusion of singleton observations in the estimations.

Table 2: Effect of 3G network availability at the time of birth on the probability of FGM

Dep var: FGM (0/1)	(1)	(2)	(3)	(4)
3G available	-0.023*** (0.005)	-0.021** (0.009)	-0.021** (0.009)	-0.021 (0.013)
N	63,760	63,760	64,018	25,775
Mean dep var.	0.169	0.169	0.169	0.169
Estimation method	TWFE	Sun & Abraham	Callaway & Sant'Anna	Chaisemartin & D'Haultfoeuille

*Note:* Columns 1–4 estimate the effect of 3G availability at birth in the cluster of residence on the probability of undergoing FGM. Column 1 reports the results of a TWFE estimation. Column 2 reports the results of the method described in Sun & Abraham (2021). Column 3 reports the results of the method described in Callaway & Sant'Anna (2021). Column 4 reports the results of the method described in Chaisemartin & D'Haultfoeuille (2020). TWFE and Sun & Abraham methods do not include all the observations in the main analytical sample presented in the descriptive statistics (64,018 observations) since singleton observations were automatically excluded. The N reported in the Chaisemartin & D'Haultfoeuille (2020) estimation includes only the effective observations used in the estimation, not the full sample of eligible observations (64,018). The mean of the dependent variable is calculated over the sample of available observations, rather than the effective observations, because the command does not retain which observations were actually used in the estimation. \*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table 3: Heterogeneous effects of 3G availability at the time of birth on the probability of FGM and placebo analyses

Dep var: FGM (0/1)	(1)	(2)	(3)	(4)	(5)	(6)	(7)
3G available	-0.019*** (0.005)	-0.005 (0.008)	-0.014** (0.007)	-0.004 (0.008)	-0.032*** (0.008)	-0.002 (0.007)	0.000 (0.011)
3G available × FGM prevalence >50%	-0.013 (0.017)						
3G available × Rural		-0.030** (0.012)					
3G available × Muslim			-0.014 (0.010)				
3G available × Main FGM benefit: avoids premarital sex				-0.139** (0.057)			
3G available × FGM linked to religion					0.031 (0.040)		
N	63,726	63,759	63,664	63,710	37,205	4,352	10,798

*Note:* Column 1 reports the heterogeneous impact of 3G availability at time of birth in the cluster of residence on the probability of having experienced FGM by whether the prevalence of FGM among women aged 15-49 is higher or lower than 50%. The prevalence of FGM in the analytical sample (girls born between 2007 and 2018) is 41.4% in areas with high prevalence of FGM and 11.1% in areas with low prevalence of FGM. Column 2 reports the heterogeneous impact of 3G availability at time of birth in the cluster of residence on the probability of having experienced FGM by whether the girl lives in a rural or urban area. The prevalence of FGM in the analytical sample is 17.7% in rural areas and 15.5% in urban areas. Column 3 reports the heterogeneous impact of 3G availability at time of birth in the cluster of residence on the probability of having experienced FGM by whether the girl lives in a Muslim family. The prevalence of FGM in the analytical sample is 24.2% among Muslims and 8.1% among non-Muslims. Column 4 reports the heterogeneous impact of 3G availability at time of birth in the cluster of residence on the probability of having experienced FGM by the share of individuals in the state in 2008 that believe preserving virginity is the main advantage for FGM. Column 5 reports the heterogeneous impact of 3G availability at time of birth in the cluster of residence on the probability of having experienced FGM by whether the family believes FGM is linked to religion. The prevalence of FGM in the analytical sample is 50.7% among daughters from women that believe FGM is linked to religion and 11.7% among daughters from women that believe FGM is not linked to religion. Column 6 reports the results of a placebo analysis estimating the impact of 3G availability at time of birth in the cluster of residence on the probability of having experienced FGM for girls that live in regions where the prevalence of FGM is significant (at least 10%) but where the preservation of virginity is not mentioned as a benefit of FGM (less than 3%). In these regions, the most cited benefits for FGM can be social acceptance, hygiene, religion, or a better marriage. Column 7 reports the results of a placebo analysis estimating the impact of 3G availability at time of birth in the cluster of residence on the probability of having experienced FGM for girls that live in regions where out-of-marriage sex is less stigmatized. We defined these regions as regions where the average number of lifetime sexual partners excluding husbands is at least 2.\*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table 4: Robustness checks: Effect of 3G availability at the time of birth on the probability of FGM

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
	FGM (0/1)	FGM (0/1)	FGM (0/1)	FGM (0/1)	FGM (0/1)	FGM (0/1)	Age at cutting	FGM (0/1)
3G available	-0.023*** (0.005)	-0.023*** (0.005)	-0.024*** (0.005)	-0.013*** (0.004)	-0.027*** (0.006)	-0.016** (0.007)	0.121 (0.091)	-0.024*** (0.007)
N	63,760	63,760	57,711	41,881	59,506	45,549	10,251	29,684
Mean dep var.	0.169	0.169	0.171	0.179	0.171	0.166	1.000	0.160
Age at survey FE	No	Yes	Yes	No	No	No	Yes	No
Control variables	No	No	Yes	No	No	No	No	No
Household FE	No	No	No	Yes	No	No	No	No
Analytical sample	All girls	All girls	All girls	All girls	Excluding potentially "partially" treated	Girls>2	Ever cut girls	Non-migrants sample

*Note:* Columns 1-6 and 8 estimate the impact of 3G availability at the time of birth in the cluster of residence on the probability of having experienced FGM. Column 1 reports the results of the main TWFE benchmark specification. Column 2 reports the results of the TWFE specification including age at survey fixed effects. Column 3 reports the results of the TWFE specification including age at survey fixed effects, and household access to electricity, TV and radio as control variables. Column 4 reports the results of the TWFE specification including household fixed effects. Column 5 reports the results of the TWFE specification excluding from the sample girls that were 1 year or 2 years old at the time of 3G arrival. These girls might be partially treated. Column 6 reports the results of the TWFE specification but excluding from the analytical sample girls younger than 3 years old at the time of the survey. More than 95% of the women cut are cut within the first 2 years of life. Column 7 reports the results of the effect of 3G on age at cutting. Column 8 reports the results of the analytical sample including only in the analytical sample those girls whose parents did not migrate during their lifetime.\*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table 5: Effect of 3G availability at the time of the survey on support for FGM

Panel A:	All		Women		Men	
Support for FGM (1-3)	(1)	(2)	(3)	(4)	(5)	(6)
Share pop with 3G available	-0.681*** (0.146)	-0.247*** (0.049)	-0.727*** (0.177)	-0.164*** (0.046)	0.319 (0.333)	0.190 (0.174)
N	39,547	83,039	30,967	61,038	8,408	21,996
Mean dep var.	1.543	1.542	1.543	1.534	1.545	1.564
Parallel trends test	0.059 (0.051)	-0.025 (0.072)	0.076 (0.081)	-0.051 (0.086)	-0.138 (0.130)	-0.028 (0.132)
Estimation method	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE
Panel B:	All		Women		Men	
Support for FGM (0/1)	(1)	(2)	(3)	(4)	(5)	(6)
Share pop with 3G available	-0.348*** (0.080)	-0.130*** (0.025)	-0.334*** (0.098)	-0.070*** (0.024)	0.124 (0.171)	0.028 (0.090)
N	39,547	83,039	30,967	61,038	8,408	21,996
Mean dep var.	0.308	0.307	0.308	0.306	0.309	0.310
Parallel trends test	0.037 (0.029)	-0.006 (0.039)	0.040 (0.045)	-0.020 (0.046)	-0.073 (0.071)	-0.018 (0.074)
Estimation method	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE

*Note:* Columns 1-6 estimate the impact of 3G availability at the time of survey on support for FGM for women aged 15-49 and men aged 15-59 surveyed in DHS rounds 2003, 2008, 2013 and 2018. In Panel A *Support for FGM* is a discrete variable that takes the value of 1 if individual does not want FGM to continue, 2 if depends, and 3 if respondent wants the practice to continue. In Panel B, *Support for FGM* is a dummy variable that takes the value of 0 if individual does not want FGM to continue, and 1 if individual responds depends or wants the practice to continue. Column 1, 3, and 5 report the results of the method described in [Chaisemartin and D'Haultfoeuille \(2020\)](#). Columns 2, 4, and 6 report the results of a two-way fixed effect model (TWFE). The test for parallel trends in the TWFE estimations is a placebo test that measures the effect of the share of population covered by 3G in 2018 on the evolution of the outcome variable between 2003 and 2008, before the expansion of the 3G network in Nigeria. The parallel trends test in the Chaisemartin & D'Haultfoeuille estimations is the placebo analysis on the pre-trends developed in [Chaisemartin and D'Haultfoeuille \(2020\)](#). TWFE estimations do not include all the observations in the main analytical sample presented in the descriptive statistics since singleton observations were automatically excluded. The N reported in the Chaisemartin & D'Haultfoeuille (2020) estimations include only the effective observations used in the estimations, not the full sample of eligible observations. The mean of the dependent variable is calculated over the sample of available observations, rather than the effective observations, because the command does not retain which observations were actually used in the estimation.\*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table 6: Effect of 3G availability at the time of the survey on sexual behavior outcomes (Chaisemartin and D’Haultfoeuille, 2020 estimation method)

	(1)	(2)	(3)	(4)	(5)	(6)
	Age at 1st marriage	Age at 1st intercourse	N lifetime sexual partners	N lifetime sexual partners excluding spouse	N sexual partners (last 12m excluding spouse)	Used condom last relation (0/1)
<i>Panel A: Women</i>						
Share pop with 3G available	0.520 (0.374)	-0.836*** (0.318)	1.028*** (0.295)	0.867*** (0.293)	0.046* (0.023)	0.014 (0.017)
N	46,719	50,454	62,254	62,254	62,582	47,406
Parallel trends test	-0.016 (0.184)	-0.079 (0.155)	-0.061 (0.057)	-0.111* (0.063)	0.006 (0.010)	-0.002 (0.008)
Mean dep var.	17.911	16.806	1.479	0.618	0.104	0.064
<i>Panel B: Men</i>						
Share pop with 3G available	1.224* (0.726)	-0.828 (0.647)	0.053 (0.653)	-0.646 (0.646)	-0.130* (0.074)	-0.053 (0.044)
N	13,222	16,675	23,296	22,448	23,774	16,108
Parallel trends test	-0.361 (0.379)	-0.182 (0.317)	-0.267 (0.232)	-0.247 (0.238)	-0.069 (0.051)	-0.028 (0.024)
Mean dep var.	25.270	20.324	3.160	2.395	0.256	0.176


*Note:* The table reports the estimates of the impact of 3G availability at time of survey on sexual behavior outcomes using the difference-in-differences method described in Chaisemartin and D’Haultfoeuille (2020). Panel A reports the estimations for women and Panel B for men. The parallel trends test is the placebo analysis on the pre-trends developed in Chaisemartin and D’Haultfoeuille (2020). The N includes only the effective observations used in the estimations, not the full sample of eligible observations. The mean of the dependent variable is calculated over the sample of available observations, rather than the effective observations, because the command does not retain which observations were actually used in the estimation. \*\*\* $p < 0.01$ ; \*\* $p < 0.05$ ; \* $p < 0.1$ .

Table 7: Effects of the anti-FGM online campaign for the International Day of the Girl Child on support for FGM

Panel A: Support for FGM (1-3)						
	(1)	(2)	(3)	(4)	(5)	(6)
Post	-0.064 (0.089)	-0.034 (0.024)	0.018 (0.067)	0.166 (0.172)	0.058 (0.040)	0.129 (0.085)
Estimation method	Non parametric	Parametric Linear fit	Parametric Quadratic fit	Non parametric	Parametric Linear fit	Parametric Quadratic fit
Cluster FE	No	Yes	Yes	No	Yes	Yes
Donut (1 week before)	No	No	No	Yes	Yes	Yes
Observations	6,484	6,482	6,482	6,106	6,102	6,102
Mean	1.392	1.392	1.392	1.384	1.384	1.384
Panel B: Support for FGM (0/1)						
	(1)	(2)	(3)	(4)	(5)	(6)
Post	-0.007 (0.052)	-0.024 (0.015)	0.027 (0.039)	-0.038 (0.093)	0.015 (0.024)	0.071 (0.050)
Estimation method	Non parametric	Parametric Linear fit	Parametric Quadratic fit	Non parametric	Parametric Linear fit	Parametric Quadratic fit
Cluster FE	No	Yes	Yes	No	Yes	Yes
Donut (1 week before)	No	No	No	Yes	Yes	Yes
Observations	6,484	6,482	6,482	6,106	6,102	6,102
Mean	0.231	0.231	0.231	0.228	0.228	0.228


*Note:* Panel A reports the estimates of the short-term effects of the anti-FGM online campaign held for the International Day of the Girl Child on October 11th, 2018, on Support for FGM. The variable Support for FGM is a discrete measure that takes the value 1 if the respondent does not want FGM to continue, 2 if the answer is “it depends,” and 3 if the respondent wants the practice to continue. In Panel B, *Support for FGM* is a dummy variable that takes the value of 0 if individual does not want FGM to continue, and 1 if individual responds depends or wants the practice to continue. Columns 1-3 report the RDD estimates for the subsample of individuals living in locations with 3G network available at the time of the campaign using the 2018 DHS data. We compute parametric estimations with polynomials of order 1 and 2 for the running variable and the non-parametric method following the procedure presented in [Calonico et al. \(2019\)](#). The parametric regressions include survey cluster fixed effects. Columns 4-6 reestimate the same regressions excluding respondents interviewed during the week before October 11th, which might be affected by the pre-campaign announcements. \*\*\* $p < 0.01$ ; \*\* $p < 0.05$ ; \* $p < 0.1$ .

Figure 1: Proportion of women subjected to FGM in Nigeria by year of birth




Note: The figure depicts the evolution of the proportion of women subjected to FGM in Nigeria by year of birth for girls and women older than 3. The figure is constructed using data from the Nigerian Demographic and Health survey, rounds 2003, 2008, 2013, and 2018, and MICS 2016, and 2021.

Figure 2: Prevalence of FGM in Nigeria by county (LGA) (age 0-49) - DHS 2018




Note: The graphs depicts the proportion of females undergoing FGM in Nigeria by county (LGA). The figure is constructed using the 2018 Nigerian DHS round, the last year for which information on 3G is available. The Nigerian DHS data is not representative at the LGA level, but at the state level. The state level prevalence is shown in [Figure A4](#).

Figure 3: Perceived benefits of FGM in Nigeria




Note: The figure shows the proportion of individuals in Nigeria that perceives FGM has each of the following benefits. Data used is the 2008 Nigerian DHS, which is the last survey including information on the perceived benefits of FGM. Male participants were aged 15–59, and female participants were aged 15–49. Participants could indicate more than one perceived benefit of FGM, as responses were not mutually exclusive.

Figure 4: Share of individuals with 3G network available by county (LGA) and year




Note: The figure illustrates the expansion of the 3G network in Nigeria using GMSA data between 2012 and 2018 and the NASA population dataset to calculate the share of population with 3G network available every year at the county level.

Figure 5: Effect of 3G network availability at the time of survey on internet access (Sun & Abraham)




Note: This figure presents event studies estimating the dynamic effects of 3G network availability at the time of the survey on internet access, using the estimator developed by Sun and Abraham (2021). The bars represent 95% confidence intervals. 3G availability is defined at the village-year level. Top graphs refer to the entire sample while bottom graphs restrict the sample to people between 18 and 30 years old. The data used are the rounds 2010, 2012, 2015 and 2018 of the Nigerian General Household Survey that surveyed the same villages over time.

Figure 6: Event study: Effect of 3G network availability at the time of birth on the probability of FGM (OLS)



Note: The graph depicts the dynamic effects of 3G availability at the time of birth on the probability of FGM using OLS methods. The bars represent 95% confidence intervals.

Figure 7: The October 11th anti-FGM online campaign and the support for FGM (1-3)



Note: The figure displays the fitted linear polynomial from a regression discontinuity design estimating the short-term effect of the anti-FGM online campaign held on October 11th, 2018, for the International Day of the Girl Child, on support for FGM (1-3), using data from the 2018 Nigerian DHS round. The plot includes 95% confidence intervals and separate fits for individuals living in areas with and without 3G network coverage.

# Appendix A Additional tables and figures

Table A1: Data sources

Outcome	Datasets	Rounds	Sample	Exposure to 3G
FGM (0/1)	DHS and MICS	2021, 2018, 2016, 2013	Daughters module	Individual exposure to 3G at year of birth
Support for FGM	DHS	2018, 2013, 2008, 2003	Female and male modules	Proportion of individuals in county exposed to 3G at time of the survey
Age at 1st marriage	DHS	2018, 2013, 2008, 2003	Female and male modules	Proportion of individuals in county exposed to 3G at time of the survey
Age at 1st intercourse	DHS	2018, 2013, 2008, 2003	Female and male modules	Proportion of individuals in county exposed to 3G at time of the survey
Lifetime sexual partners	DHS	2018, 2013, 2008, 2003	Female and male modules	Proportion of individuals in county exposed to 3G at time of the survey
Sexual partners excluding spouse	DHS	2018, 2013, 2008, 2003	Female and male modules	Proportion of individuals in county exposed to 3G at time of the survey
Used condom last relation	DHS	2018, 2013, 2008, 2003	Female and male modules	Proportion of individuals in county exposed to 3G at time of the survey
Labor force partic.	DHS	2018, 2013, 2008, 2003	Female and male modules	Proportion of individuals in county exposed to 3G at time of the survey
Hh wealth index	DHS	2018, 2013, 2008, 2003	Female and male modules	Proportion of individuals in county exposed to 3G at time of the survey
Internet access	NGHS	2018, 2015, 2012, 2010	Female and male modules	Individual exposure to 3G at the time of the survey

*Note:* Table provides information on the data sources used in the analysis of the effect of 3G on different outcomes, as well as on how exposure to 3G is defined.

Table A2: Robustness check: Anti-FGM legislation

Dep. var.: FGM (0/1)	(1)	(2)	(3)
3G available	-0.023*** (0.005)	-0.024*** (0.005)	-0.024*** (0.006)
N	63,760	63,760	51,171
Mean dep. var.	0.169	0.169	0.176
Estimation method	TWFE	TWFE	TWFE
Control variable: FGM ban at birth	No	Yes	No
Analytical sample	All girls	All girls	Excluding regions with anti-FGM laws

*Note:* Columns 1–3 estimate the impact of 3G availability at the time of birth in the cluster of residence on the probability of having experienced FGM. All columns use TWFE estimation. Column 1 reports the benchmark result from column (1) of Table 2. Column 2 adds a control for whether a law banning FGM was in place in the region at the time of the girl’s birth. Column 3 excludes girls living in regions that had already adopted legislation banning FGM by 2018, the last year for which we have information on 3G availability.\*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table A3: Placebo test: Effect of 2G availability at the time of birth on the probability of FGM

Dep var: FGM (0/1)	(1)	(2)	(3)	(4)
2G available	0.010* (0.005)	0.005 (0.008)	0.023 (0.020)	-0.005 (0.018)
N	39,929	39,929	26,502	18,998
Mean dep var.	0.181	0.181	0.176	0.175
Estimation method	TWFE	Sun & Abraham	Callaway & Sant'Anna	Chaisemartin & D'Haultfoeuille

*Note:* Columns 1–4 estimate the effect of 2G availability at birth in the cluster of residence on the probability of undergoing FGM. Column 1 reports the results of a TWFE estimation. Column 2 reports the results of the method described in Sun & Abraham (2021). Column 3 reports the results of the method described in Callaway & Sant'Anna (2021). Columns 4 reports the results of the method described in Chaisemartin & D'Haultfoeuille (2020).\*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table A4: Effect of 3G availability on state presence: Public schools openings, access to electricity, radio and TV

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
	N schools opened	N schools opened	Access to electricity (0/1)	Access to electricity (0/1)	Radio (0/1)	Radio (0/1)	TV (0/1)	TV (0/1)
Share pop with 3G available	-0.503 (0.399)	0.157 (0.171)	-0.115 (0.095)	0.015 (0.029)	0.057 (0.054)	0.068*** (0.021)	-0.038 (0.063)	-0.047* (0.025)
N	5,110	8,760	86,489	169,054	86,489	169,078	86,489	169,063
Mean dep var.	1.379	1.379	0.524	0.528	0.703	0.704	0.461	0.465
Parallel trends test	0.007 (0.116)	0.267 (0.258)	-0.029 (0.044)	0.109* (0.056)	0.009 (0.020)	-0.029 (0.030)	-0.010 (0.029)	-0.025 (0.056)
Estimation method	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE

*Note:* Columns 1–8 estimate the impact of 3G availability on different measures of state presence. Columns 1–2 show the effect of 3G availability in the LGA on the opening of public schools. Columns 3–4 show the effect of 3G availability at the time of the survey on access to electricity. Columns 5–6 show the effect of 3G availability at the time of the survey on having a radio. Columns 7–8 show the effect of 3G availability at the time of the survey on having a TV. Columns 1, 3, 5, and 7 report the results of the method described in [Chaisemartin and D'Haultfoeuille \(2020\)](#), while Columns 2, 4, 6, and 8 report the results of a two-way fixed effects model (TWFE). The test for parallel trends in the TWFE estimations is a placebo test that measures the effect of the share of population covered by 3G in 2018 on the evolution of the outcome variable before the start of the deployment of 3G network in Nigeria. The parallel trends test in the [Chaisemartin and D'Haultfoeuille \(2020\)](#) estimations is the placebo analysis on pre-trends proposed by the authors. In the [Chaisemartin and D'Haultfoeuille \(2020\)](#) estimations, the mean of the dependent variable is calculated over the sample of available observations, rather than the effective sample, because the command does not retain which observations were actually used in the estimation.\*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table A5: Effect of 3G availability at the time of the survey on support for FGM (including the 2016 MICS round)

Panel A:	All		Women		Men	
Dep var: Support for FGM (1-3)	(1)	(2)	(3)	(4)	(5)	(6)
Share pop with 3G available	-0.540*** (0.125)	-0.183*** (0.041)	-0.522*** (0.149)	-0.086** (0.039)	0.319 (0.333)	0.190 (0.174)
N	52,845	100,181	44,237	78,181	8,408	21,996
Mean dep var.	1.533	1.535	1.533	1.527	1.545	1.564
Parallel trends test (p-value)	0.370	0.731	0.307	0.551	0.291	0.836
Estimation method	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE
Panel B:	All		Women		Men	
Dep var: Support for FGM (0/1)	(1)	(2)	(3)	(4)	(5)	(6)
Share pop with 3G available	-0.258*** (0.066)	-0.098*** (0.021)	-0.215*** (0.079)	-0.034* (0.020)	0.124 (0.171)	0.028 (0.090)
N	52,845	100,181	44,237	78,181	8,408	21,996
Mean dep var.	0.302	0.303	0.302	0.301	0.309	0.310
Parallel trends test (p-value)	0.275	0.872	0.211	0.668	0.304	0.806
Estimation method	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE

*Note:* Columns 1-6 estimate the impact of 3G availability at the time of survey on support for FGM for women aged 15-49 and men aged 15-59 surveyed in DHS rounds 2003, 2008, 2013 and 2018, and MICS 2016. In Panel A *Support for FGM* is a discrete variable that takes the value of 1 if individual does not want FGM to continue, 2 if depends, and 3 if respondent wants the practice to continue. In Panel B, *Support for FGM* is a dummy variable that takes the value of 0 if individual does not want FGM to continue, and 1 if individual responds depends or wants the practice to continue. Column 1, 3, and 5 report the results of the method described in Chaisemartin and D'Haultfoeuille (2020). Columns 2, 4, and 6 report the results of a two-way fixed effect model (TWFE). The test for parallel trends is a placebo test that measures the effect of the share of population covered by 3G in 2018 on the evolution of the outcome variable between 2003 and 2008, before the expansion of the 3G network in Nigeria. The parallel trends test in the Chaisemartin & D'Haultfoeuille estimations is the placebo analysis on the pre-trends developed in Chaisemartin and D'Haultfoeuille (2020). TWFE estimations do not include all the observations in the main analytical sample presented in the descriptive statistics since singleton observations were automatically excluded. The N reported in the Chaisemartin & D'Haultfoeuille (2020) estimations include only the effective observations used in the estimations, not the full sample of eligible observations. The mean of the dependent variable is calculated over the sample of available observations, rather than the effective observations, because the command does not retain which observations were actually used in the estimation. \*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table A6: Effect of 3G availability at the time of the survey on sexual behavior outcomes (TWFE estimation method)

	(1)	(2)	(3)	(4)	(5)	(6)
	Age at 1st marriage	Age at 1st intercourse	N lifetime sexual partners	N lifetime sexual partners excluding spouses	N sexual partners (last 12m excluding spouses)	Used condom last relation (0/1)
<i>Panel A: Women</i>						
Share pop with 3G available	0.564*** (0.156)	-0.105 (0.138)	0.577** (0.262)	0.515* (0.267)	0.015 (0.012)	-0.015 (0.009)
N	90,609	97,002	112,213	112,106	120,587	91,717
Parallel trends test	0.054 (0.329)	0.167 (0.200)	NA	NA	0.001 (0.019)	0.039** (0.016)
Mean dep var.	17.938	16.822	1.480	0.621	0.104	0.064
<i>Panel B: Men</i>						
Share pop with 3G available	0.140 (0.273)	-0.164 (0.304)	-0.273 (0.301)	-0.444 (0.299)	-0.077*** (0.026)	-0.081*** (0.020)
N	27,181	34,670	44,397	41,633	47,943	32,929
Parallel trends test	0.160 (0.653)	0.153 (0.467)	NA	NA	0.233** (0.116)	0.006 (0.042)
Mean dep var.	25.286	20.323	3.174	2.409	0.256	0.177

*Note:* The table reports the estimates of the impact of 3G availability at time of survey on sexual behavior outcomes using TWFE. Panel A reports the estimations for women and Panel B for men. TWFE estimations do not include all the observations in the main analytical sample presented in the descriptive statistics since singleton observations were automatically excluded. The test for parallel trends in the TWFE estimations is a placebo test that measures the effect of the share of population covered by 3G in 2018 on the evolution of the outcome variable between 2003 and 2008, before the expansion of the 3G network in Nigeria. Since information on the number of lifetime sexual partners is not included in the 2003 DHS round, the pre-trends test in the TWFE specification cannot be estimated for these variables. \*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table A7: Effect of 3G availability at the time of the survey on sexual behavior outcomes for individuals aged 30 or older (Chaisemartin and D’Haultfoeuille, 2020 estimation method)

	(1)	(2)	(3)	(4)	(5)	(6)
	Age at 1st marriage	Age at 1st intercourse	N lifetime sexual partners	N lifetime sexual partners excluding spouse	N sexual partners (last 12m excluding spouse)	Used condom last relation (0/1)
<i>Panel A: Women</i>						
Share pop with 3G available	0.642 (0.464)	-0.878** (0.388)	1.305*** (0.406)	1.231*** (0.405)	0.075*** (0.023)	0.019 (0.014)
N	27,211	26,530	27,961	27,961	28,221	24,677
Parallel trends test	-0.076 (0.240)	-0.076 (0.223)	-0.079 (0.076)	(0.086)	-0.131 -0.012 (0.010)	0.002 (0.010)
Mean dep var.	18.670	17.077	1.844	0.706	0.050	0.034
<i>Panel B: Men</i>						
Share pop with 3G available	0.762 (0.788)	-1.637** (0.768)	-1.036 (1.030)	-1.562 (1.115)	-0.093 (0.076)	-0.003 (0.036)
N	11,047	11,345	11,696	10,874	12,116	11,362
Parallel trends test	-0.411 (0.451)	-0.185 (0.426)	-0.143 (0.464)	-0.226 (0.539)	-0.018 (0.044)	-0.015 (0.019)
Mean dep var.	26.037	21.264	4.491	3.321	0.171	0.099

*Note:* The table reports the estimates of the impact of 3G availability at time of survey on sexual behavior outcomes for individuals aged 30 or older using the difference-in-differences method described in Chaisemartin and D’Haultfoeuille (2020). Panel A reports the estimations for women and Panel B for men. The N includes only the effective observations used in the estimations, not the full sample of eligible observations. The parallel trends test is the placebo analysis on the pre-trends developed in Chaisemartin and D’Haultfoeuille (2020). The mean of the dependent variable is calculated over the sample of available observations, rather than the effective observations, because the command does not retain which observations were actually used in the estimation.\*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table A8: Effect of 3G availability at the time of the survey on wealth and labor force participation

	(1)	(2)	(3)	(4)
	HH wealth index	HH wealth index	Labor force participation	Labor force participation
Panel A: All indiv.				
Share pop with 3G availability	-0.041 (0.143)	0.000 (0.049)	0.022 (0.048)	0.006 (0.016)
N	86,489	169,172	86,257	168,568
Parallel trends test	-0.028 (0.064)	0.007 (0.099)	-0.018 (0.021)	0.032 (0.034)
Mean dep var.	0.000	0.000	0.672	0.672
Estimation method	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE
Panel B: Women				
Share pop with 3G availability	-0.053 (0.144)	-0.002 (0.049)	-0.027 (0.063)	-0.010 (0.020)
N	62,651	120,998	62,485	120,516
Parallel trends test	-0.023 (0.064)	0.014 (0.098)	-0.031 (0.026)	0.070 (0.043)
Mean dep var.	-0.018	-0.018	0.616	0.616
Estimation method	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE
Panel C: Men				
Share pop with 3G availability	-0.017 (0.150)	0.001 (0.051)	0.114** (0.046)	0.038* (0.021)
N	23,838	48,173	23,772	48,051
Parallel trends test	-0.023 (0.064)	0.001 (0.121)	-0.031 (0.026)	-0.037 (0.039)
Mean dep var.	0.044	0.044	0.811	0.811
Estimation method	Chaisemartin & D'Haultfoeuille	TWFE	Chaisemartin & D'Haultfoeuille	TWFE

*Note:* Columns 1-4 report the estimates of the impact of 3G availability at the time of the survey on household wealth index and on labor force participation. The wealth index varies between -284155 and 315067 but for estimation purposes the variable has been normalized. TWFE estimations do not include all the observations in the main analytical sample presented in the descriptive statistics since singleton observations were automatically excluded. The N reported in the Chaisemartin & D'Haultfoeuille (2020) estimations include only the effective observations used in the estimations, not the full sample of eligible observations. The test for parallel trends in the TWFE estimations is a placebo test that measures the effect of the share of population covered by 3G in 2018 on the evolution of the outcome variable between 2003 and 2008, before the expansion of the 3G network in Nigeria. The parallel trends test in the Chaisemartin & D'Haultfoeuille estimations is the placebo analysis on the pre-trends developed in Chaisemartin and D'Haultfoeuille (2020). The mean of the dependent variable is calculated over the sample of available observations, rather than the effective observations, because the command does not retain which observations were actually used in the estimation. \*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table A9: Effect of 3G availability at the time of the survey on bargaining power of women within the household

	(1)	(2)	(3)	(4)	(5)
Respondent	Women	Women	Women	Women	Men
Final say on	Health care	Large HH purchases	Visits family relatives	Manage women money	Large HH purchases
<i>Panel A: Chaisemartin &amp; D'Haultfoeuille</i>					
Share pop with 3G available	0.171* (0.088)	0.135 (0.083)	0.044 (0.102)	0.081 (0.075)	-0.237 (0.149)
N	43,445	43,423	43,448	42,846	12,841
Parallel trends test	0.044 (0.046)	0.003 (0.036)	-0.015 (0.044)	-0.005 (0.052)	0.017 (0.049)
Mean dep var.	1.497	1.440	1.640	1.342	2.571
<i>Panel B: TWFE</i>					
Share pop with 3G available	0.153*** (0.038)	0.117*** (0.034)	0.089** (0.039)	-0.023 (0.037)	-0.284*** (0.063)
N	84,930	84,860	84,912	78,326	26,652
Parallel trends test	-0.105 (0.088)	-0.147*** (0.053)	0.059 (0.075)	0.000 (0.000)	0.011 (0.049)
Mean dep var.	1.502	1.446	1.648	1.344	2.507


*Note:* The table reports the estimates of the impact of 3G availability at the time of the survey on bargaining power within the household for men and women. The variables indicate the final say of the respondent on different household decisions for married individuals. The variable takes the value 3 if individual takes the decision alone, 2 if individuals participate in the decision, and 1 if the decision is taken by another person. Columns (1) to (4) report the results for women and column (5) reports the results for men. Panel A reports the estimations of the difference-in-differences method described in [Chaisemartin and D'Haultfoeuille \(2020\)](#). Panel B reports the results of a two-way fixed effects model. The test for parallel trends in the TWFE estimations is a placebo test that measures the effect of the share of population covered by 3G in 2018 on the evolution of the outcome variable between 2003 and 2008, before the expansion of the 3G network in Nigeria. The parallel trends test in the Chaisemartin & D'Haultfoeuille estimations is the placebo analysis on the pre-trends developed in [Chaisemartin and D'Haultfoeuille \(2020\)](#). The event study is reported in [Figure A22](#). TWFE methods do not include all the observations in the main analytical sample presented in the descriptive statistics since singleton observations were automatically excluded. The N reported in the Chaisemartin & D'Haultfoeuille (2020) estimations include only the effective observations used in the estimations, not the full sample of eligible observations. The mean of the dependent variable is calculated over the sample of available observations, rather than the effective observations, because the command does not retain which observations were actually used in the estimation.\*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table A10: Effect of 3G availability on schooling outcomes

	(1) Educ. Attainment (0-4)	(2) Secondary educ (0/1)	(3) School enrollment (0/1)	(4) School enrollment (0/1)
Panel A: All indiv.				
N years exposed to 3G during schooling years (0-13)	-0.020*** (0.004)	-0.004* (0.002)		
Share pop with 3G available			-0.122 (0.078)	-0.022 (0.023)
N	88,226	88,226	93,832	181,980
Parallel trends test			-0.005 (0.038)	0.108*** (0.036)
Mean dep var.	1.374	0.534	0.680	0.680
Estimation method	OLS	OLS	Chaisemartin & D'Haultfoeuille	TWFE
Panel B: Women				
N years exposed to 3G during schooling years (0-13)	-0.018*** (0.005)	-0.002 (0.002)		
Share pop with 3G available			-0.138* (0.082)	-0.036 (0.024)
N	64,629	64,629	46,361	90,025
Parallel trends test			0.009 (0.038)	0.091** (0.037)
Mean dep var.	1.268	0.490	0.651	0.651
Estimation method	OLS	OLS	Chaisemartin & D'Haultfoeuille	TWFE
Panel C: Men				
N years exposed to 3G during schooling years (0-13)	-0.024*** (0.007)	-0.006* (0.003)		
Share pop with 3G available			-0.109 (0.081)	-0.007 (0.024)
N	23,575	23,575	47,466	91,945
Parallel trends test			-0.018 (0.041)	0.128*** (0.041)
Mean dep var.	1.661	0.653	0.709	0.709
Estimation method	OLS	OLS	Chaisemartin & D'Haultfoeuille	TWFE


*Note:* Columns 1-4 report the estimates of the effect of exposure to 3G on different schooling outcomes. Column 1 reports the OLS estimates of the number of years of exposure to 3G during the school years (ages 6 to 18) on educational attainment for individuals aged 18 or older. Educational attainment is a discrete variable that takes the value 0 if the individual has no education, 1 if primary education, 2 if secondary education, and 3 if higher education. Column 2 reports the OLS estimates of the number of years of exposure to 3G during the school years (ages 6 to 18) on the probability of having secondary education for individuals aged 18 or older. Columns 3 and 4 report the difference-in-differences estimates (see equation 2) of exposure to 3G at the time of the survey on school enrollment at the time of the survey for individuals aged 6 to 18. Column 3 reports estimates using the procedure developed by Chaisemartin and D'Haultfoeuille, and Column 4 reports results from the canonical TWFE. The test for parallel trends in the TWFE estimations in column 3 is a placebo test that measures the effect of the share of population covered by 3G in 2018 on the evolution of the outcome variable between 2003 and 2008, before the expansion of the 3G network in Nigeria. The parallel trends test in the Chaisemartin & D'Haultfoeuille estimations in column 4 is the placebo analysis on the pre-trends developed in [Chaisemartin and D'Haultfoeuille \(2020\)](#).\*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Figure A1: Prevalence of FGM by country and age group




Note: The figure shows the average prevalence of FGM across two groups of women: Women aged 15-19, and women aged 45-49, for those countries with an average prevalence of FGM larger than 10% among women aged 15-49. Because the vast majority of FGM occurs by the age of 15, the differences in the prevalence of FGM across generations captures the variation in the practice over time. To build the figure, we use data from the last round of DHS and MICS surveys with information on FGM in every African country. An exception to this is Nigeria. While a MICS survey was conducted in 2021, we use the 2018 DHS because it is the last year for which information on 3G coverage is available and thus the last year in the studied period. The descriptive statistics in the rest of the figures are therefore based on the 2018 DHS, and for consistency, we also use the same dataset here. However, the trends in the figure do not change relevantly when the Nigerian MICS 2021 data are used.

Figure A2: Support for FGM by country and sex




Note: The figure shows the average support for FGM by sex across different countries, focusing on men and women aged 15–49. Surveys typically include nationally representative samples of men aged 15–59 and women aged 15–49. Since our interest is in comparing support across genders, we restrict the male sample to ages 15–49, excluding men aged 50–59. Support for FGM is defined as the percentage of respondents who answer “yes” or “depends” to the question of whether the practice should continue. The data are drawn from the most recent DHS or MICS survey available for each country that includes information on support for FGM by sex.

Figure A3: Prevalence of FGM in Nigeria by county (LGA) (age 0-49) - DHS 2008




Note: The graphs depicts the proportion of females undergoing FGM in Nigeria by county. The figure is constructed using the 2008 Nigerian DHS round, the last round before the deployment of 3G. The Nigerian DHS data is not representative at the county level, but at the regional level. The regional level prevalence is shown in Figure A4.

Figure A4: Prevalence of FGM in Nigeria by region (age 15-49)




Note: The graphs depict the proportion of females aged 15-49 subjected to FGM in Nigeria by state. The figure is constructed using the 2018 and 2008 Nigerian DHS round. These surveys were representative at the regional level for this population group.

Figure A5: Internet access in Nigeria




Note: The graph shows the evolution of the proportion of the population that report having access to internet in Nigeria. The proportions are calculated using the Nigerian General Household Survey rounds 2010, 2012, 2015, and 2018.

Figure A6: Main form of internet access in Nigeria over time




Note: The figure shows the evolution of the main sources of internet access in Nigeria. The share for each source is calculated over the sample of individuals with internet access. The data come from the 2010, 2012, 2015, and 2018 rounds of the Nigerian General Household Survey.

Figure A7: Effect of 3G availability at the time of the survey on internet access (OLS)




Note: This figure presents event studies estimating the dynamic effects of 3G availability at the time of the survey on internet access using OLS. The bars represent 95% confidence intervals. 3G availability is defined at the village-year level. Top graphs refer to the entire sample while bottom graphs restrict the sample to people between 18 and 30 years old. The data used are the rounds 2010, 2012, 2015 and 2018 of the Nigerian General Household Survey that surveyed the same villages over time.

Figure A8: Event study: Effect of 3G availability at the time of birth on the probability of FGM (Sun & Abraham)




Note: The figure depicts the dynamic effects of 3G availability at the time of birth on the probability of FGM using the Sun & Abraham event study. The bars represent 95% confidence intervals.

Figure A9: Effect of 2G network availability at the time of birth on the probability of FGM




Note: The figure shows the dynamic effects of 2G network availability at the time of birth on the probability of FGM using the event study estimator presented in Sun and Abraham (2021). The bars represent 95% confidence intervals.


Figure A10: Effect of 3G availability at the time of the survey on different measures of state presence




(a) N of schools opened



(b) Access to electricity




(c) Has a radio



(d) Has a TV


Note: The figure shows the dynamic effects of 3G network availability at the time of the survey on different measures of state presence using the event study estimator developed in [Chaisemartin and D'Haultfœuille \(2020\)](#). The bars represent 95% confidence intervals. For the analysis of electricity, radio, and TV, we use data from the 2010, 2012, 2015, and 2018 rounds of the DHS. For the analysis of public school openings, we construct a yearly panel of school openings at the county level between 2007 and 2018.

Figure A11: Effect of 3G network availability at the time of the survey on the support for FGM (1-3)



(a) All individuals


(b) Women



(c) Men


Note: The figure shows the dynamic effects of 3G network availability at the time of the survey on the support for FGM using the event study estimator developed in [Chaisemartin and D'Haultfoeuille \(2020\)](#). 3G availability is defined as the share of the population in a county (LGA) covered by 3G. Data used include rounds 2003, 2008, 2013, and 2018 of the Nigerian Demographic and Health Surveys. Information on support for FGM is not included for men in DHS 2018. The bars represent 95% confidence intervals.

Figure A12: Effect of 3G network availability at the time of the survey on the support for FGM (0/1)



(a) All individuals

(b) Women




(c) Men


Note: The figure shows the dynamic effects of 3G network availability at the time of the survey on the support for FGM (0/1) using the event study estimator developed in [Chaisemartin and D'Haultfoeulle \(2020\)](#). 3G availability is defined as the share of the population in a county (LGA) covered by 3G. Data used include rounds 2003, 2008, 2013, and 2018 of the Nigerian Demographic and Health Surveys. Information on support for FGM is not included for men in DHS 2018. The bars represent 95% confidence intervals.

H


Figure A13: Effect of 3G network availability at the time of the survey on sexual behavior outcomes for women




(a) Age at 1st marriage




(b) Age at 1st intercourse




(c) N lifetime sexual partners



(d) N lifetime sexual partners (excluding spouses)



(e) N extramarital partners last 12 months




(f) Used condom in last relation


Note: The figure shows the dynamic effects of 3G network availability at the time of the survey on sexual behavior outcomes for women using the estimator presented in [Chaisemartin and D'Haultfoeuille \(2020\)](#). 3G availability is defined as the share of the population in a county (LGA) covered by 3G. Data used include the Nigerian Demographic and Health Surveys, rounds 2003, 2008, 2013, and 2018. The bars represent 95% confidence intervals.

## H


Figure A14: Effect of 3G network availability at the time of the survey on sexual behavior outcomes for men




(a) Age at 1st marriage




(b) Age at 1st intercourse




(c) N lifetime sexual partners



(d) N lifetime sexual partners (excluding spouses)



(e) N extramarital partners last 12 months



(f) Used condom in last relation


Note: The figure shows the dynamic effects of 3G network availability at the time of the survey on sexual behavior outcomes for men using the estimator presented in [Chaisemartin and D'Haultfœuille \(2020\)](#). 3G availability is defined as the share of the population in a county (LGA) covered by 3G. Data used include the Nigerian Demographic and Health Surveys, rounds 2003, 2008, 2013, and 2018. The bars represent 95% confidence intervals.

Figure A15: Examples of anti-FGM tweets




Note: The figure displays various examples of anti-FGM tweets in Nigeria. The tweets were issued on October 11th, 2018, during the International Day of the Girl Child.

Figure A16: Daily Number of anti-FGM tweets in 2018




Note: The figure depicts the daily number of tweets in Nigeria that used one or more of the following anti-FGM hashtags: #endFGM, #endcuttinggirls, or #nofgm. The two vertical lines on February 6th and October 11th indicate the International Day of Zero Tolerance for FGM and the International Day of the Girl Child, respectively. No survey fieldwork was conducted in 2018, nor in any other year during the days immediately before or after the International Day of Zero Tolerance for FGM (Feb 6th).

Figure A17: Google searches for FGM in Nigeria



Note: Google trends data. The figure depicts Google searches between 2014 and 2021 for the main words used to refer to FGM in Nigeria in English, Yoruba, Igbo, and Hausa, the main languages spoken in Nigeria. The searches are normalized to the moment and term of more Google searches.


Figure A18: The October 11th anti-FGM online campaign and the support for FGM (0/1)



The figure displays the fitted linear polynomial from a regression discontinuity design estimating the short-term effect of the anti-FGM online campaign held on October 11th, 2018, for the International Day of the Girl Child, on support for FGM (0/1), using data from the 2018 Nigerian DHS round. The plot includes 95% confidence intervals and separate fits for individuals living in areas with and without 3G network coverage.

## H


Figure A19: Effect of 3G network availability at the time of the survey on household wealth



Note: The figure shows the dynamic effects of 3G network availability at the time of the survey on household wealth using the estimator presented in [Chaisemartin and D'Haultfoeuille \(2020\)](#). 3G availability is defined as the share of the population in a county (LGA) covered by 3G. Data used include the Nigerian Demographic and Health Surveys, rounds 2003, 2008, 2013, and 2018. The bars represent 95% confidence intervals.



## H

Figure A20: Effect of 3G network availability at the time of the survey on labor force participation



Note: The figure shows the dynamic effects of 3G network availability at the time of the survey on labor force participation using the estimator presented in [Chaisemartin and D'Haultfoeuille \(2020\)](#). 3G availability is defined as the share of the population in a county (LGA) covered by 3G. Data used include the Nigerian Demographic and Health Surveys, rounds 2003, 2008, 2013, and 2018. The bars represent 95% confidence intervals.


Figure A21: Effect of 3G availability at the time of the survey on school enrollment (individuals aged 6-18)



Note: The figure shows the dynamic effects of 3G network availability at the time of the survey on school enrollment for individuals aged 6-18 using the event study estimator developed in [Chaisemartin and D'Haultfœuille \(2020\)](#). 3G availability is defined as the share of the population in a county (LGA) covered by 3G. Data used include rounds 2003, 2008, 2013, and 2018 of the Nigerian Demographic and Health Surveys. The bars represent 95% confidence intervals.

## H

Figure A22: Effect of 3G network availability at the time of the survey on final say on...



Note: The figure shows the dynamic effects of 3G network availability at the time of the survey on bargaining power within the household for women and men using the estimator presented in [Chaisemartin and D'Haultfoeulle \(2020\)](#). 3G availability is defined as the share of the population in a county (LGA) covered by 3G. Data used include the Nigerian Demographic and Health Surveys, rounds 2003, 2008, 2013, and 2018. The bars represent 95% confidence intervals.

## Appendix B 3G network availability, support for FGM, and sexual behavior: 3G availability defined at the individual level

In this appendix, we explore the association between 3G network availability, support for FGM, sexual behavior and other outcomes using an alternative empirical approach. Rather than using the share of individuals in the county (LGA) with 3G network available at the time of the survey, we define an alternative measure of treatment exposure based on whether the individual lives in a village with 3G network available at the time of the survey. Specifically, we estimate the following regression:

$$\begin{aligned} Outcome_{it} = & \delta_0 + \delta_1 3G\ network\ available_{it} + \delta_2 Never\ received\ 3G\ network_i \\ & + Year\ of\ interview_t + \nu_l + u_{ilt} \end{aligned} \quad (5)$$

where  $Outcome_{it}$  indicates the outcome of interest for individual  $i$ , living in county (LGA)  $l$  and interviewed in year  $t$ .  $3G\ network\ available_{it}$  is a dummy variable that indicates whether individual  $i$  interviewed in year  $t$  has 3G network available.  $Never\ received\ 3G\ network$  is a dummy variable equal to 1 if individual  $i$  lives in a location that never received 3G during the study period.  $Year\ of\ interview_{it}$  and  $\nu_l$  are year-of-interview (survey) and county fixed effects.  $u$  is the error term. Standard errors are clustered by county. The main parameter of interest is  $\delta_1$ , which measures the association between the availability of 3G network and the outcome of interest.

Unlike in the main analysis, where exposure to 3G is measured as the share of individuals in the county with 3G network availability, exposure to 3G networks in this analysis is defined at the individual level based on the geocoded location of villages or neighborhoods. However, this approach is problematic because individuals and villages in the sample are only interviewed once during the study period. Thus, the empirical framework used is not a difference-in-differences and the causal interpretation of the analysis does not rely on the parallel trends assumption. Rather, the interpretation of  $\delta_1$  as the causal effect of 3G network availability on the outcome of interest mainly relies on no unobservable variables correlated with both timing of 3G network expansion and the outcome of interest. To limit endogeneity concerns, we introduced a variable indicating whether individual  $i$  lives in a location unexposed during the study period, which implies that the variation used for the identification of the parameter of interest  $\delta_1$  comes from individuals living in locations that at

some point during the study period received 3G network, but that were interviewed either before or after the arrival of the network.

The results of these estimations broadly support the main conclusions of the study. The estimates for FGM support are reported in Table B1, showing negative associations between availability of 3G networks and both measures of the support for FGM, which are mainly driven by women.

Also in line with the evidence provided above, the results reported in Table B2 show that the availability of 3G networks is associated with a higher age at marriage and a higher number of lifetime sexual partners for women, and a lower age at first intercourse for men. Like in the main analysis, there is no association between 3G availability and wealth or labor force participation for both men or women.

Table B1: 3G network availability at the time of the survey and support for FGM: Individual level exposure to 3G network

	All		Women		Men	
	Support for FGM (1-3)	Support for FGM (0/1)	Support for FGM (1-3)	Support for FGM (0/1)	Support for FGM (1-3)	Support for FGM (0/1)
	(1)	(2)	(3)	(4)	(5)	(6)
3G available	-0.095*** ( 0.029)	-0.051*** ( 0.015)	-0.057** ( 0.027)	-0.024 ( 0.014)	0.020 ( 0.090)	-0.010 ( 0.048)
N	83,039	83,039	61,038	61,038	21,996	21,996
Mean dep var.	1.542	0.307	1.534	0.306	1.564	0.310

*Note:* Columns 1-6 estimate the association using OLS between 3G network availability at the time of the survey measured at the individual level and the support for FGM among women aged 15-49 and men aged 15-59 surveyed in DHS rounds 2003, 2008, 2013 and 2018. In Panel A *Support for FGM* is a discrete variable that takes the value of 1 if individual does not want FGM to continue, 2 if depends, and 3 if respondent wants the practice to continue. In Panel B, *Support for FGM* is a dummy variable that takes the value of 0 if individual does not want FGM to continue, and 1 if individual responds depends or wants the practice to continue. The regressions estimated include LGA fixed effects, year of interview fixed effects, and a dummy variable indicating whether the village of residence did not receive 3G during the study period. Thus, the variation used in the estimations arises from locations that at some point receive 3G network during the study period but were interviewed either before or after the arrival of 3G network. Columns 1 and 2 report the results using the full sample, Columns 3 and 4 report the results using the sample of women, and Columns 5 and 6 report the results using the male sample. Standard errors are clustered at the LGA level. \*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

Table B2: 3G network availability at the time of the survey and sexual behavior outcomes, labor force participation, and household wealth: Individual level exposure to 3G network

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
	Age at 1st marriage	Age at 1st intercourse	N lifetime sexual partners	N lifetime sexual partners (excluding husb.)	N sexual partners (last 12m excluding husb.)	Used condom last relation (0/1)	Labor force participation	HH wealth
<i>Panel A: Women</i>								
3G available	0.262** ( 0.103)	-0.000 ( 0.082)	0.244* ( 0.133)	0.205 ( 0.136)	-0.000 ( 0.008)	-0.011** ( 0.005)	0.001 ( 0.012)	0.009 ( 0.033)
N	90,609	97,002	112,213	112,106	120,587	91,717	120,516	120,998
Mean dep var.	17.938	16.822	1.480	0.621	0.104	0.064	0.616	-0.018
<i>Panel B: Men</i>								
3G available	-0.101 ( 0.192)	-0.285* ( 0.166)	-0.062 ( 0.180)	-0.149 ( 0.178)	-0.028 ( 0.019)	-0.048*** ( 0.014)	0.016 ( 0.013)	-0.009 ( 0.037)
N	27,181	34,670	44,397	41,633	47,943	32,929	48,051	48,173
Mean dep var.	25.286	20.323	3.174	2.409	0.256	0.177	0.811	0.044

*Note:* Columns 1-8 estimate the association using OLS between 3G network availability at time of survey measured at the individual level and sexual behavioral outcomes, household wealth, and labor force participation for women aged 15-49 and men aged 15-59 surveyed in DHS rounds 2003, 2008, 2013 and 2018. The regressions estimated include LGA fixed effect, year of interview fixed effects, and a dummy variable indicating whether the village of residence did not receive internet during the study period. Thus, the variation used in the estimations arises from locations that at some point receive 3G network during the study period but were interviewed either before or after the arrival of 3G network. Panel A reports the results using the sample of women, and Panel B reports the results using the male sample. Standard errors are clustered at the LGA level.\*\*\*p<0.01; \*\*p<0.05; \*p<0.1.

## Appendix C What internet content is normalizing premarital sex?

This appendix explores which specific types of internet content may be driving the change in sexual behavior and contributing to the erosion of premarital sex stigma. To address this question, we conduct the following analyses. First, we conduct focus groups in 8 rural communities in the areas of Calabar and Abuja, gathering the views from 68 Nigerian women from all major ethnic groups in Nigeria. Second, to shed further light on the results of the qualitative research, we use Google searches to provide a broader picture about internet content consumed in Nigeria. Finally, we revisit the existing literature on the effects of internet content on sexual behavior and discuss the feasibility of the qualitative findings.

### C.1 Qualitative research

In December 2024, we conducted 10 focus groups throughout 8 rural communities, gathering the views from 68 Nigerian women in rural areas of Abuja and Calabar. Additionally, we conducted in-depth interviews with 6 Christian and Muslim religious and traditional leaders. The communities surveyed in the area of Calabar

were Christian, predominantly from Efok, Ekoi, and Ibibio ethnicities. In the area of Abuja, we visited Igbo, Yoruba and Hausa rural communities. Individuals living in these rural communities near Abuja were eminently internal migrants that moved from Igboland (east), Yorubaland (south west), and the North of the country in search for better economic opportunities and security in the last few years.

Each focus group lasted for approximately 1 hour and a half, and while fluent English was spoken by most of the participants, the research team—two of the study authors—was supported by a local translator that also facilitated the focus groups. The discussions were mainly focused on internet use and their perceived effects of internet on social norms, particularly on sexual behaviors and stigmas.

Participants in the focus groups emphasized three main internet uses. First, communication with friends and family living outside the community. Second, they reported using internet for entertainment such as films, music, and social media. While the specific content varied across age groups, both younger and older women appeared to use the internet for similar purposes, though the consumption of Western cultural content was more prevalent among the younger participants. Third, most participants also use internet to access updated information, particularly news media and weather forecasts. Finally, some focus groups also highlighted educational and business-related uses of the internet among some participants.

And what were the perceived effects of internet on sexual behavior? Despite coming from a wide range of ethnic groups throughout the country, the women interviewed offered remarkably similar accounts about how the expansion of internet access—mostly through smartphones since the early 2010s—has reshaped social norms, particularly around premarital sex. Once widely stigmatized and associated with shame or a loss of family honor, premarital sex is now increasingly seen as a normal part of youth life, especially among younger generations. In Christian communities, it was consistently described as “the new normal,” noting that young people face little, if any, social sanction from families or traditional leaders. This stands in sharp contrast to a few years ago, when such behavior could damage a girl’s marriage prospects or trigger strong parental discipline. In Muslim communities, while premarital sex is reportedly on the rise among young people as well, it remains a sensitive and taboo topic for religious leaders. This rapid change is generating tensions in some Muslim communities, particularly with traditional and religious leaders who continue to strongly oppose such practices.

When asked what specific online content was triggering the shift in sexual norms and behaviors, the women interviewed highlighted two types of content. The first includes films, music, and social media posts depicting Western cultural content

in which premarital sex is portrayed as normal. The second consists of films and other content that depict nudity or explicit sexual acts. Many emphasized that “young people do everything they see online”—not just in terms of sexual behavior, but also in how they dress, the music they listen to, and how they interact with others. Respondents described this exposure as a key factor triggering the observed changes in behavior, by introducing new role models and reshaping perceptions of what is normal or acceptable. While television was occasionally mentioned, it was consistently described as less influential due to shared viewing spaces and greater family oversight. In contrast, mobile phones provide private, unregulated access to global content, which respondents identified as central to the speed and depth of these transformations—even in communities that, until recently, were relatively isolated.

Additionally, although religious and traditional leaders continue to denounce premarital sex—particularly in Muslim communities—their influence appears to have weakened. In some cases, interviewees pointed to inconsistencies between leaders’ messages and their own behavior. As a result, religious discourse remains present but is perceived as less effective in regulating sexual behavior.

Beyond sexual norms, internet access has brought both opportunities and concerns. On the positive side, it facilitates communication—especially with migrants—, access to information, and educational or commercial activities. Some respondents praised its value for business, learning, and staying connected with the wider world. At the same time, internet use is frequently associated with moral decline. Terms like “corruption” or “copying bad behavior” appeared in nearly all discussions, typically referring to sexual content, fraudulent schemes, and a general departure from local customs and values.

In multiple interviews, internet access was also linked to rising rates of teenage pregnancy and out-of-marriage pregnancies. While poverty and parental neglect were also cited as contributing factors, many participants stressed that exposure to sexual content online encourages early sexual activity. Interestingly, a few women in Christian communities noted that among educated young women, internet access could also have protective effects by increasing knowledge about contraception.

## **C.2 Google trends analysis**

While the results of the qualitative research presented above emphasized the powerful influence of both Western cultural content and sexually explicit content in shaping young people’s behavior, it remains important to ask whether access to such content is, in fact, widespread in Nigeria. To explore this question, we exam-

ined the 200 most searched terms on Google in Nigeria.<sup>50</sup> Among these we identified some expected terms related to news, social media, films, popular national media, and UK and Spanish football. We also found several terms that may help understand how Nigerians from traditional communities are exposed to Western cultural content, such as YouTube, Instagram, Facebook, and websites providing free access to films. Moreover, among the 200 most cited terms, we identified 11 pornographic websites, which together account for more than 14.6 million searches per month. This figure refers only to Google searches—actual traffic to websites is likely to be significantly higher, as many users access them directly or through other platforms.

Figure C1: 20 most searched terms in Google

KEYWORD	VOLUME	flashscore mobi	4,090,000
livescore	9,140,000	xnxx	4,090,000
spartybet	7,480,000	livescore cz	3,350,000
translate	6,120,000	betking	3,350,000
flashscores	6,120,000	whether	2,740,000
bet9ja shop	5,000,000	weather	2,740,000
bet9ja	5,000,000	porn	2,240,000
bbc hausa	5,000,000	youtube	2,240,000
facebook	4,090,000	man united	1,830,000

Note: The figure depicts the most 20 searched terms and the approximate volume of monthly searches in Nigeria in 2024.

### C.3 Discussion and previous evidence on the effects of exposure to Western cultural content and sexually explicit content

The vast majority of women interviewed identified two important triggers of the normalization of premarital sex: (1) Western cultural content in which premarital sex is portrayed as normal, and (2) sexually explicit content. While we cannot test the specific effects of these types of content in our setting, is there any evidence on the effects of exposure to these contents?

Regarding the effect of exposure to Western cultural content, a few studies show that exposure to Western media can transform beliefs and behaviors, bringing them more in line with the values portrayed in these media. [Jensen and Oster \(2009\)](#)

<sup>50</sup>Figure C1 shows the 20 most searched terms and the approximate volume of monthly searches in Nigeria.

shows that access to cable TV in India reduced the acceptability of wife beating, lowered the reported preference for sons, increased female school enrollment, and decreased fertility. [La Ferrara et al. \(2012\)](#) shows that soap operas portraying small families led to lower fertility. However, media can also backfire: [Blumenstock et al. \(2022\)](#) shows that, in Pakistan, exposure to Western media increased voting for religious parties and enrollment in religious schools.

A growing body of literature indicates that exposure to sexually explicit content is associated with shifts in sexual norms and behaviors, including a greater acceptance of premarital sex and earlier sexual initiation. A systematic review by [Pathmendra et al. \(2023\)](#) synthesizes the existing evidence and concludes that higher levels of pornography exposure are consistently associated with a younger age at first sexual intercourse. Supporting this pattern, [Brown et al. \(2006\)](#) find that exposure to sexual content accelerates adolescents' sexual activity and increases their risk of engaging in early sexual intercourse. Beyond behavioral outcomes, [Peter and Valkenburg \(2008\)](#) show that pornography consumption has been linked to heightened sexual preoccupation and [Vandenbosch and Eggermont \(2015\)](#) document that exposure to pornographic websites is directly associated with the internalization of appearance ideals, self-objectification, and increased body surveillance, highlighting the broader influence of pornography on adolescents' sexual self-concept and body image.

But why exposure to sexually explicit content might affect sexual behaviors? Different mechanisms have been proposed in the literature. First, pornography often serves as a form of informal sex education, shaping young people's beliefs about what is typical or acceptable sexual behavior. Second, sexually explicit content may encourage behavioral imitation (or copy-cutting), as adolescents replicate the sexual acts, roles, or dynamics they observe, assuming these portrayals reflect real-life expectations. Third, exposure to sexually explicit content can lower perceived social and psychological barriers to engaging in sex by framing it as casual, risk-free, and universally practiced. Fourth, repeated exposure may lead to desensitization, weakening internalized moral or cultural norms and shifting perceptions of peer behavior. Finally, sexually explicit content can elicit sexual arousal, which may heighten short-term motivation to initiate sexual activity, particularly in contexts where opportunities are available.

While our qualitative results show that exposure to sexually explicit content is perceived by the local population as a key contributor to the normalization and destigmatization of premarital sex—with implications for practices such as female genital cutting—our aim is to explain observed patterns, not to make normative

judgments about this content. We neither endorse its use nor propose it as a policy tool, and we do not overlook the potential harms associated with it. A substantial body of research has documented associations between pornography consumption and a range of adverse outcomes, including increased sexual aggression ([Waterman et al., 2022](#); [Wright et al., 2016](#)), aggressive attitudes and behaviors ([Zhou et al., 2021](#)), and poorer mental health indicators ([Setyawati et al., 2020](#)). Although causality remains difficult to establish due to methodological constraints, these findings highlight legitimate concerns about the possible negative consequences of exposure to sexually explicit content, particularly during adolescence.

## Appendix D The effect of rainfall shocks on FGM in Nigeria

One potential mechanism driving the effect of 3G network on FGM is an increase in income. If 3G networks increase household income (Bahia et al., 2024) and FGM increases a daughter’s value in marriage (i.e., leading to a higher bride price) (Khalifa, 2022; García-Hombrados and Salgado, 2022), parents at the margin of cutting their daughters might decide to stop the practice because the additional income from 3G compensates for the loss of income associated with not having their daughter subjected to FGM. The latter mechanism implies that 3G might decrease the prevalence of FGM even if it does not reduce the support for this practice.

To test this hypothesis, we examine in this appendix the effect of a pure income shock on FGM, specifically rainfall shocks. If a pure income shock does not affect the prevalence of FGM, it is unlikely that the effect of 3G on FGM is solely driven by a rise in economic welfare. To this end, we replicate the analysis conducted by McGavock and Novak (2023), focusing on Nigeria. The latter paper assesses the effect of rainfall shocks during the age of risk of FGM on the prevalence of this practice in sub-Saharan Africa. Following McGavock and Novak (2023), we estimate the following equation using Nigerian data:

$$FGM_{ibv} = \delta_0 + \delta_1 Rainfall\ shock_{bv} + \theta_b + \gamma_v + u_{ibv} \quad (6)$$

where *Rainfall shock* is equal to 1 if the annual precipitation in cluster  $v$  during the year of birth  $b$  was below a certain percentile of the gamma distribution of the historical rainfall in the cluster.<sup>51</sup> Because the typical age at cutting varies across sub-Saharan African countries, McGavock and Novak (2023) focuses on rainfall shocks during the year the girl reached the first percentile of the distribution of age at FGM. Given that in Nigeria, more than 80% of the cuts occur within the first year of life, we focus on rainfall shocks during the year of birth.  $\theta_b$  and  $\gamma_v$  are survey cluster and year of birth fixed effects.  $u$  is the error term.

The results of this analysis are reported in columns 1–4 of Table D1, using different percentiles of the historical rainfall distribution within the village to define negative economic shocks. Our findings show that the estimated coefficients of rainfall shocks across these different definitions are negative and mostly not statistically different from zero. These results suggest that, if anything, negative economic

---

<sup>51</sup>The rainfall information was provided by Lindsey Novak and Tamara McGavock, who constructed a database on annual precipitation in DHS clusters between 1900 and 2017 using monthly precipitation information gathered by the University of Delaware at the 0.5 x 0.5 degree grid.

shocks reduce the prevalence of FGM, supporting the hypothesis that positive economic shocks are unlikely to be a major driver of the effects of 3G on FGM. Our results align with those of [McGavock and Novak \(2023\)](#), who find that, in some settings, negative economic shocks increase FGM prevalence, but observe no effect for ethnic groups where cutting occurs early in life, as in the case of Nigeria.

As an additional empirical exercise, we re-estimate equation 1 including rainfall shocks as an additional control variable to assess whether meteorological shocks could confound the effect of 3G availability. Column 5 in Table D1 shows the benchmark estimation of the effect of 3G availability at the time of birth reported in Table 2, while column 6 reports the same estimation including rainfall shocks as a control variable. The estimates are virtually identical, reassuring that rainfall shocks are not confounding the main results of the study.

Table D1: Effect of adverse rainfall shocks at the time of birth on the probability of FGM in Nigeria

Dep var: FGM (0/1)	(1)	(2)	(3)	(4)	(5)	(6)
Adverse rainfall shock	-0.002 (0.012)	-0.011 (0.007)	-0.012* (0.006)	-0.003 (0.006)		
3G available					-0.023*** (0.005)	-0.023*** (0.005)
Rainfall percentile defining drought	Lowest 5%	Lowest 10%	Lowest 15%	Lowest 20%		
Rainfall shock control					No	Yes
N	63,760	63760	63,760	63,760	63,760	63,760
Mean dep. var	0.169	0.169	0.169	0.169	0.169	0.169

*Note:* Columns 1–4 report the estimated impact of rainfall shocks at the time of birth on the probability of female genital mutilation (FGM). Rainfall shocks are defined as years in which rainfall falls within the lowest 5%, 10%, 15%, and 20% of the historical distribution of rainfall within a cluster, corresponding to columns 1 through 4 respectively. Column 5 reports the benchmark estimation of the effect of 3G availability at the time of birth on FGM. Column 6 reports the main estimation of the effect of 3G availability at the time of birth on the probability of FGM, including rainfall shocks as a control variable. Rainfall shocks in this specification are defined as years in which rainfall falls within the lowest 15% of the historical distribution of rainfall within each cluster. \*\*\*p<0.01; \*\*p<0.05; \*p<0.1.